arXiv:1212.2618v1 [math.GR] 11 Dec 2012

# SURFACE SUBGROUPS FROM LINEAR PROGRAMMING

DANNY CALEGARI AND ALDEN WALKER

Abstract. We show that a *random endomorphism* of a free group gives rise to an HNN extension which contains a closed surface subgroup, with probability one; a special case is the HNN extension associated to the endomorphism of a rank 2 free group (considered by Sapir) sending $a$ to $ab$ and $b$ to $ba$.

We further show that a group obtained by doubling a free group along any collection of subgroups (of arbitrary rank) has the property that every rational 2-dimensional homology class is virtually represented by surface subgroups, and the unit ball in the Gromov norm is a finite sided rational polyhedron.

These results are obtained by a mixture of combinatorial, geometric and linear programming techniques. We obtain other related results of a more technical nature concerning stable commutator length in free groups, especially concerning the structure of the space of so-called *counting quasimorphisms*

## Contents

1. Introduction 1
2. Stable commutator length 5
3. Traintracks 8
4. Counting functions and posimorphisms 12
5. The geometry of $Q_\ell$ 19
6. Stallings traintracks and surface subgroups of doubles 26
7. Endomorphisms and traintrack dynamics 28
8. Folded surfaces and $f$-folded surfaces 34
9. Extremal surfaces and flat surfaces 48
10. Acknowledgments 56
References 56

## 1. Introduction

1.1. **Gromov's surface subgroup question.** The following well-known question is usually attributed to Gromov:

**Question 1.1** (Gromov). *Let $G$ be a one-ended hyperbolic group. Does $G$ contain the fundamental group of a closed surface with $\chi < 0$?*

Hereafter we abbreviate "fundamental group of a closed surface with $\chi < 0$" to "surface group", so that this question asks whether every one-ended hyperbolic group contains a surface subgroup. This question is wide open in general, but a positive answer is known in certain special cases, including:

---

*Date*: December 12, 2012.

1. Coxeter groups (Gordon–Long–Reid [21]);
2. Graphs of free groups with cyclic edge groups and $b_2 > 0$ (Calegari [7]);
3. Fundamental groups of hyperbolic 3-manifolds (Kahn–Markovic [27]);
4. Certain doubles of free groups (Kim–Wilton, Kim–Oum [29, 28]);

(this list is not exhaustive).

One of the goals of this paper is to describe how linear programming may be used to settle the question of the existence of surface subgroups in certain graphs of free groups, either by giving a powerful computational tool to *find* surface subgroups in specific groups, or by reducing the analysis of this question in infinite families of groups to a finite (tractable) calculation.

We are able to prove the existence of surface subgroups in the following groups:

1. A group $G$ with $b_2 > 0$ obtained by doubling a free group $F$ along a finite collection of finitely generated subgroups $F_i$ (see § 6.3);
2. A group $G$ obtained as an HNN extension $F*_\phi$ where $F$ is a free group of fixed rank and $\phi$ is a *random* endomorphism (see § 8.3);
3. "Sapir's group" $C = F*_\phi$ for $F = \langle a, b\rangle$ and $\phi : a \to ab, b \to ba$ (see § 8.4).

One sense in which this constitutes a significant advance over the results in [7, 29, 28] (for instance) is that the edge groups are free groups of *arbitrary rank*, whereas in the cited papers the edge groups were required to be cyclic.

Bullet (1) above is implied by a stronger result about the Gromov norm on $H_2$ of the double of $F$ along the $F_i$; we discuss this stronger result in the next subsection. Bullet (3) is reasonably self-explanatory. A precise statement of bullet (2) is:

**Random $f$-folded Surface Theorem 8.9.** *Let $k \geq 2$ be fixed, and let $F$ be a free group of rank $k$. Let $\phi$ be a random endomorphism of $F$ of length $n$. Then the probability that $F*_\phi$ contains an essential surface subgroup is at least $1 - O(C^{-n^c})$ for some $C > 1$ and $c > 0$.*

Here a random endomorphism of length $n$ is one that takes the generators to reduced words of length $n$ chosen independently and randomly with the uniform distribution. We became interested in surface subgroups of HNN extensions of free groups after discussions with Mark Sapir, who conjectured that the subgroup $C$ does not contain a surface subgroup, and thought it was unlikely that many HNN extensions should contain surface subgroups (other than $\mathbb{Z}^2$ subgroups for endomorphisms fixing a nontrivial conjugacy class). Therefore it seems safe to say that the Random $f$-folded Surface Theorem is in many ways very unexpected.

The reader who is interested only in Gromov's surface subgroup question can read § 8 with minimal reference to the rest of the paper. § 6 depends more substantially on the material in § 3.

### 1.2. Stable commutator length and Gromov norm.

We are interested not only in the *existence* of surface subgroups, but also in their *quality*. Fix a group $G$. Every integral 2-dimensional homology class (resp. relative 2-dimensional homology class) in $G$ is represented by a nontrivial — but not necessarily injective — map from a surface subgroup (resp. surface with boundary subgroup). The complexity of such a surface is measured by $-\chi/2$, and the infimum of the complexity of the surfaces projectively representing a given homology class is $1/4$ times the *Gromov norm* of the class (resp. the scl *norm* of the relative class). A surface realizing the infimum may not exist, but if it does, the map is *injective*. We distinguish such special surface subgroups from more general ones by calling them *extremal*.

If $G$ is a graph of free groups, we can find a surface subgroup of $G$ as an amalgam of surface subgroups with boundary in the free factors. Thus one way to find an extremal subgroup of $G$ is to build it out of subgroups with boundary extremal in the factors. To specify a 2-dimensional homology class necessitates the choice of a homologically trivial family $\Gamma$ of conjugacy classes in $G$. Then $H_2(G,\Gamma)$ is an affine space over $H_2(G)$ (for instance, if $H_2(G)$ is trivial, $H_2(G,\Gamma)$ consists of a single class). If $G$ is a graph of free groups amalgamated over cyclic subgroups, there are natural candidates for $\Gamma$, namely (powers of) the cyclic generators of the edge groups. But for amalgams over higher rank groups, there are infinitely many conjugacy classes to consider, even up to taking powers.

For $F$ a free group and $\Gamma$ a specific homologically trivial family of conjugacy classes, the scl norm of the nontrivial class in $H_2(G,\Gamma)$ is just the (usual) stable commutator length of $\Gamma$, and denoted $\text{scl}(\Gamma)$ (see [8] for an introduction to this subject). In [9] an algorithm is given to compute $\text{scl}(\Gamma)$ in a free group, and the algorithm proves that the infimum is realized by an extremal surface. One of the main results in this paper is an extension of this algorithm to minimize $\text{scl}(\Gamma)$ over certain *infinite* families $\Gamma$; roughly speaking, those parameterized by *weights on a traintrack*.

The main theorem we prove along these lines is the *Traintrack Rationality Theorem*:

**Traintrack Rationality Theorem 3.24.** *For $w \in W^+$, there is a formula*

$$\text{scl}(w) = \inf\{-\chi(y)/2 \ : \ y \in Y^+ \text{ and } D_E(y) = w\}$$

*Moreover, the infimum is achieved, and if $w$ is rational, the infimum is achieved at a rational $y$. Furthermore, the function* $\text{scl}(w)$ *varies piecewise rational linearly on $W^+$.*

Here $W^+$ denotes the non-negative homologically trivial weights on a traintrack over a rose for $F$ (for precise definitions, see § 3).

A double of such an extremal surface with boundary is extremal in the double, so we obtain the following theorem:

**Double Norm Theorem 6.5.** *Let $F$ be a free group, and $F_i$ a finite collection of finitely generated subgroups. Let $G$ be the double of $F$ along the $F_i$; i.e. $G := D(F, \{F_i\})$. Then the unit ball in the Gromov norm on $H_2(G)$ is a rational polyhedron, and every rational class in $H_2(G)$ is projectively represented by a finite sum of injective closed surface subgroups.*

The same methods allow one to approximate in a natural way the (polyhedral) unit ball in the scl norm in the infinite dimensional space $B_1^H(F)$ for $F$ a free group by a natural sequence of (polyhedral) unit balls in *finite dimensional* normed spaces. The definition of these finite dimensional spaces, and the computation of the norms, is entirely effective, and we describe an explicit and efficient algorithm to do so.

### 1.3. **Counting functions and combinatorics.**

Stable commutator length is a quantitative version of relative 2-dimensional homology. It is dual to 2-dimensional *bounded cohomology*, as introduced by Gromov [24]. More precisely, for any group $G$ there is an exact sequence

$$0 \to H^1(G) \to Q(G) \to H_b^2(G) \to H^2(G)$$

where $Q$ denotes the vector space of *homogeneous quasimorphisms*, i.e. functions $\phi : G \to \mathbb{R}$ satisfying $\phi(g^n) = n\phi(g)$, and $|\phi(gh) - \phi(g) - \phi(h)| \le D(\phi)$ for some least constant $D(\phi)$ (called the *defect*), for any $g, h \in G$. The defect defines a norm on $Q/H^1$, and this normed vector space is the isometric dual of $B_1^H(G)$ with (twice) its scl (pseudo)-norm.

Even in a free group, our understanding of $Q$ is far from complete. Topologically, $Q(F)/H^1(F)$ is a nonseparable Banach space, and it is probably hopeless to give an explicit description of its geometry. However one may aim to give an explicit description of a subset of $Q$ large enough to witness the geometry of $B_1^H$ by operator duality. That is to say: given $b \in B_1^H$, we would like to be able to construct some explicit $\phi \in Q$ with $\text{scl}(b) = \phi(b)/2D(\phi)$; such a $\phi$ is said to be *extremal* for $b$.

A natural class of explicit homogeneous quasimorphisms are the so-called *counting quasimorphisms*, first introduced by Rhemtulla [32], and then further discussed by Brooks [5]. For a word $\sigma$, let $C^\sigma(b)$ count the number of copies of $\sigma$ appearing in $b$. A counting quasimorphism is a linear combination of functions of the form $C^\sigma - C^{\sigma^{-1}}$. If we bound the length of the words $\sigma$ appearing in such a linear combination by $\ell$, we get a space $Q_\ell$ of homomgeneous quasimorphisms. In § 4 we construct an explicit family of finite dimensional vector spaces $W_\ell/\epsilon$ with norms $\|\cdot\|_\ell$, and a compatible family of linear projections $\Psi_\ell : B_1^H \to W_\ell$ for which the following duality holds:

**Counting Quasimorphism Duality Theorem 4.24.** *For any $\ell$ and any $b \in B/\epsilon$, there is an equality*

$$\|\Psi_\ell(b)\|_\ell = \sup_{\phi \in Q_\ell/H^1} \frac{\phi(b)}{2D(\phi)}$$

It is reasonable to conjecture that $\text{scl}(b) = \lim_{\ell\to\infty} \|\Psi_\ell(b)\|_\ell$, and in fact that equality is achieved for all sufficiently large $\ell$.

By breaking the asymmetry of the $Q_\ell$, and considering the broader class of *all* counting functions (i.e. linear combinations of $C^\sigma$ as above) we are able to detect the full geometry of the scl norm ball. This depends on the notion of *posimorphisms*, which are related to quasimorphisms roughly as subharmonic functions are to harmonic functions; see Definition 4.11, or Theorem 4.19 for the analog of the Counting Quasimorphism Duality Theorem for (counting) posimorphisms.

The unit balls in $Q_\ell/H^1$ are rational polyhedra. In § 5 we go some way towards giving a description of these polyhedra. First, for $\ell = 2$ and arbitrary rank, we are able to give a complete description of the codimension one faces. Fix $S$ a symmetric free generating set for $F$, so if $F$ has rank $k$, the set $S$ has $2k$ elements. Each cyclically ordered triple of distinct elements of $F$ determines a codimension 1 face of $Q_2/H^1$, called a *tripod face*. The unit ball has the following description:

**Unit Ball Theorem 5.11.** *The codimension 1 faces of the unit ball in $Q_2/H^1$ are exactly the set of length 1 tripod faces. The symmetric group $S_{2k}$ acts on this set by permutation of edge labels, and the resulting action on $Q_2/H^1$ is by isometries. As a representation of $S_{2k}$, the space $Q_2/H^1$ is isomorphic to $\wedge^2 V$ where $V$ is the standard $(2k-1)$-dimensional representation.*

The simplicity of the statement of the Unit Ball Theorem obscures how surprising it really is; only the subgroup $\mathbb{Z}/2\mathbb{Z}^k \rtimes S_k$ of $S_{2k}$ acts on $S$ by automorphisms. The fact that the full group $S_{2k}$ of permutations acts even *linearly* on $Q_2/H^1$, let alone by isometries, is very surprising and entirely unexpected.

The generalization to bigger $\ell$ is less complete. It turns out that there are natural analogs of tripod faces in $Q_\ell$, and these are a complete set of supporting hyperplanes for the unit ball. However for $\ell \geq 3$ it is a *redundant* collection, and it is more difficult to give a finite description of the unit ball analogous to the $\ell = 2$ case. On the other hand, we are able to give a conditional answer to the problem of finding an explicit extremal quasimorphism dual to a given chain $\Gamma$. We are able to show (Corollary 5.6) that for any $b \in B_1^H$, the (possibly empty) set of $\phi \in Q_\ell/H^1$ extremal for $\Gamma$ can be computed explicitly and directly from the combinatorics of any extremal surface for $\Gamma$. In words: an extremal surface has as a spine a certain kind of *fatgraph*; this fatgraph determines a collection of tripods, namely the tripods which *immerse* into the fatgraph. The $\phi \in Q_\ell/H^1$ extremal for $\Gamma$ are then precisely the cone on the intersection of the associated tripod faces (if nonempty). For more details, see § 5.2.

1.4. **Geometric endomorphisms, and flat surfaces.** In § 9 we investigate the class of *extremal* and *flat* surfaces in HNN extensions of free groups. An extremal surface is one which realizes the Gromov norm of some class in $H_2(F*_\phi)$. We interpret the (Gromov) norm of this class as the *translation length* of $\phi$ on a certain (infinite dimensional) affine normed space modeled on $B/\epsilon$. Moreover, for *random* endomorphisms $\phi$, lower bounds for this norm can be obtained by considering the translation length on *finite dimensional* normed spaces modeled on the various $W_\ell$. If equality holds, extremal surfaces exist, and can be certified by counting quasimorphisms.

These affine actions factor through a semidirect product of the form $\mathbb{R}^N \rtimes G$ for *finite* groups $G$; we thereby obtain homomorphisms from certain subsemigroups of $\mathrm{End}(F)$ to (abelian) groups of *translations* on finite dimensional vector spaces. This can be thought of in a natural way as an analog of (a summand of) the Johnson homomorphism on the Torelli subgroup of a mapping class group. In the particular special case of *flat surfaces* the analogy is very tight.

1.5. **The role of linear programming.** What is the role of linear programming in all this? The essence of our method depends on the fact that injectivity in hyperbolic groups can often be certified by *local* data. Thus building a surface subgroup in a hyperbolic group reduces to many simple "subproblems". These problems are typically interrelated in a complicated (but also local) way, but sometimes different tools (combinatorics, ergodic theory) can be used to reduce a large collection of "weakly coupled" subproblems to a large collection of *independent* subproblems, parameterized by *linear* data in the form of linear equalities and inequalities. This last system can be solved with linear programming (hereafter abbreviated to LP).

The power and sophistication of modern practical algorithms for LP problems thus frequently lets us reduce a theoretical question concerning infinitely many groups to a concrete LP problem, and then to answer it. Note that the finite LP problem to which the theoretical question reduces might itself be quite substantial. For an example, see the proof of the Random $f$-folded Surface Theorem in § 8, especially the discussion in § 8.3.6.

## 2. Stable commutator length

In this section we recall some basic elements from the theory of stable commutator length, especially in free groups. For a reference, see [8], especially Chapter 2.

The main novelty in this section is our treatment of homogenization. We are careful to define homogenization (of group chains, 1-manifolds or surfaces) to be the equivalence relation which equates taking an (oriented) $n$-fold cover with multiplication by $n$ for *positive* $n$.

Throughout this section and the remainder of the paper, we use the notation $\mathbb{R}[*]$ for the $\mathbb{R}$-vector space with basis a given generating set $*$, and by $\mathbb{R}^+[*]$ the positive orthant. This is an $\mathbb{R}$-vector space with a canonical basis, so it makes sense to talk about integral/rational/positive vectors. We also use the notation $\mathbb{Z}[*]$, $\mathbb{Z}^+[*]$ and so on.

2.1. **Chains and antisymmetry.** We fix a free group $F$ and a free generating set $a, b, c, \cdots$. We denote inverse by capitalization, hence $A := a^{-1}$ and so on. Let $S = \{a, A, b, B, \cdots\}$ denote the symmetrized generating set. For simplicity, we work over $\mathbb{R}$ in what follows. Most of what we say makes sense for an arbitrary group $G$, but it is convenient to focus on the case of free groups.

The proof of the following lemma is obvious:

**Lemma 2.1.** *Conjugacy classes in $F$ are in bijection with reduced cyclic words in the generators.*

We let $C$ denote the free $\mathbb{R}$-vector space generated by primitive conjugacy classes in $F$, and let $C^+$ denote the positive orthant. There is a *homogenization* map from $\mathbb{R}^+[F] \to C^+$ sending $g^n \to ng$ for $n$ positive, and this map extends by linearity to $\mathbb{R}[F] \to C$.

There is an abelianization map $\text{ab} : C \to H_1$, and its restriction to $C^+$ is surjective.

**Definition 2.2.** Define $\epsilon : C \to C$ to be the involution defined on conjugacy classes by $\epsilon : g \to -g^{-1}$.

**Lemma 2.3.** *The map* ab *factors through $\epsilon$, and there is a surjection of* $\mathbb{R}^+$ *modules* $C^+ \to C/\epsilon$.

*Proof.* Since $a = -A$ in $H_1$ and so on, the first statement is obvious. The second statement is just the observation that every element of $C/\epsilon$ has an expression as a formal sum with all coefficients positive. ☐

*Remark* 2.4. We call an element of $C^+$ *primitive* if it does not contain a term of the form $t(c + c^{-1})$ with $t > 0$. Every element of $C/\epsilon$ has a unique expression as a primitive element of $C^+$.

**Definition 2.5.** We let $B$ denote the kernel of ab, so that there is an exact sequence $B \to C \to H_1$. Define $B^+ := B \cap C^+$.

**Lemma 2.6.** *$B^+$ spans $B$ as a vector space, and $B^+ \to B/\epsilon$ is a surjection of* $\mathbb{R}^+$ *modules.*

*Proof.* Write $b \in B$ as $b^+ - b^-$ with $b^+$ and $b^-$ in $C^+$. Then $b = (b^+ - \epsilon(b^-)) - (b^- - \epsilon(b^-))$ and both of the bracketed expressions are in $B^+$. The second statement follows from Lemma 2.3 and the fact that ab factors through $\epsilon$. ☐

It may seem clumsy to work with $\mathbb{R}^+$ modules, but in fact this is crucial for what follows. Note that although $C^+$ has a natural basis as a free $\mathbb{R}^+$ module, namely the set of all reduced cyclic words in the generators, the space $C/\epsilon$, which is canonically isomorphic to the set of primitive elements in $C^+$, does *not* have a natural basis as an $\mathbb{R}$ vector space. Such a basis amounts to a choice of one element for each $\epsilon$ orbit. We will return to this issue in the sequel.

2.2. **Spaces.** We fix $X$, a $K(F,1)$. For instance, it is convenient to take $X$ to be a wedge of $|S|/2$ circles, one for each mutually inverse pair of generators.

Conjugacy classes in $F$ are in bijection with free homotopy classes of maps $S^1 \to X$.

**Definition 2.7.** Let $\mathcal{C}$ denote the space of $\mathbb{R}$ linear formal sums of free homotopy classes of maps $S^1 \to X$ modulo the relation that if $\pi : L' \to L$ is an oriented cover of oriented 1-manifolds of (positive) degree $n$, then the composition $L' \to L \to X$ represents $n$ times $L \to X$.

Informally, we refer to $\mathcal{C}$ as the space of *multicurves*. With this definition, there is an obvious isomorphism of $C$ with $\mathcal{C}$ and $C^+$ with the positive orthant $\mathcal{C}^+$. Similarly, $B$ and $B^+$ are identified with the *homologically trivial* multicurves. There is a linear map $\epsilon : \mathcal{C} \to \mathcal{C}$ which reverses orientation and multiplies coefficients by $-1$. Evidently there is a natural identification of $C/\epsilon$ with $\mathcal{C}/\epsilon$, and therefore a map $\mathrm{ab} : \mathcal{C}/\epsilon \to H_1$.

A surface $\Sigma$ is *admissible* if it is compact and oriented, and if each component has non-positive Euler characteristic. A map $\Sigma \to X$ is admissible if $\Sigma$ is admissible.

*Remark* 2.8. It is convenient to allow some components of $\partial\Sigma$ to map homotopically trivially to $X$; we will refer to such components later as “ideal components”; however, they do not play a significant role in what follows.

**Definition 2.9.** We let $\mathcal{S}$ denote the space of $\mathbb{R}$ linear formal sums of free homotopy classes of admissible maps $\Sigma \to X$ modulo the relation that if $\pi : \Sigma' \to \Sigma$ is an oriented cover of (positive) degree $n$, then the composition $\Sigma' \to \Sigma \to X$ represents $n$ times $\Sigma \to X$.

Similarly, we define $\mathcal{S}^+$ as the positive orthant, and $\epsilon : \mathcal{S} \to \mathcal{S}$ reverses orientation and multiplies coefficients by $-1$. Again, $\mathcal{S}^+ \to \mathcal{S}/\epsilon$ is an isomorphism of $\mathbb{R}^+$ modules.

**Definition 2.10.** If $\Sigma$ is admissible, and $\Sigma \to X$ is a free homotopy class of map, the restriction of the map to $\partial\Sigma$ represents a class in $\mathcal{C}^+$. This defines a linear map $\partial : \mathcal{S} \to \mathcal{C}$ which restricts to $\partial : \mathcal{S}^+ \to \mathcal{C}^+$ and descends to $\partial : \mathcal{S}/\epsilon \to \mathcal{C}/\epsilon$.

*Remark* 2.11. Note that the image $\partial\mathcal{S}/\epsilon$ consists of precisely the kernel of ab, and can therefore be identified with $B/\epsilon$.

If $\chi$ denotes Euler characteristic, then because $\chi$ is multiplicative under covers, there is a well-defined linear function $\chi$ on $\mathcal{S}^+$. The function $-\chi/2$ is non-negative on $\mathcal{S}^+$, and invariant under $-\epsilon$, and therefore defines a pseudo-norm on $\mathcal{S}/\epsilon$ (thought of as an $\mathbb{R}$ vector space). With this definition in place, we have the following proposition:

**Proposition 2.12.** *Let $b \in B/\epsilon$, and think of $b$ as an element of $\mathcal{C}/\epsilon$ as above. Then*

$$\mathrm{scl}(b) = \inf_{\partial s = b} -\chi(s)/2$$

*where the infimum is taken over all $s \in \mathcal{S}/\epsilon$.*

The reader can take this as a definition of scl. For a proof (in which scl is given its usual, algebraic definition) see [8], Prop. 2.74.

*Remark* 2.13. The vector space $B/\epsilon$ is denoted $B_1^H$ in [8] and elsewhere.

## 3. TRAINTRACKS

In this section we define traintracks, and prove the Traintrack Rationality Theorem; i.e. Theorem 3.24. This theorem is proved by encoding the computation of scl as a rational linear programming problem; the linear programming problem depends rational linearly on the traintrack weight. This proves the Traintrack Rationality Theorem.

### 3.1. Traintracks.

We use the notation $\iota$ for the involution on $S$ that interchanges a generator with its inverse. At the level of homology, $\iota = -\epsilon$.

**Definition 3.1.** A *traintrack* $T$ is a directed graph with edges $E$ and vertices $V$, together with an *evaluation* function $h : E \to S$. There are *source* and *target* maps $s : E \to V$ and $t : E \to V$ giving the structure of a directed graph. An ordered pair of edges $e_0, e_1$ is *composable* if $te_0 = se_1$. A traintrack is *reduced* if for every composable pair $(e_0, e_1)$ we have $he_0 \neq \iota he_1$.

If $F$ and $S$ need to be stressed, we talk about a traintrack *over* $F, S$.

*Remark* 3.2. Note that our traintracks are *oriented*; by contrast, unoriented (and even unorientable) traintracks are important in the Thurston theory of surface automorphisms.

Recall that $X$, our $K(F, 1)$, is a bouquet of circles, one for each pair of generators for $F$. If we think of $T$ as an (oriented) 1-complex, then $h$ is just a simplicial map $T \to X$.

**Definition 3.3.** Given a traintrack with edges $E$ and vertices $V$, define linear maps $\partial : \mathbb{R}[E] \to \mathbb{R}[V]$ on generators by $\partial(e) = te - se$ and $H : \mathbb{R}[E] \to H_1$ by extending $h$ on generators by $\mathbb{R}$ linearity. Define the *weight space* $W$ to be the kernel

$$W := \ker \partial \oplus H : \mathbb{R}[E] \to \mathbb{R}[V] \oplus H_1$$

We define the *cone of positive weights* $W^+ := W \cap \mathbb{R}^+[E]$.

**Lemma 3.4.** *$W^+$ is a convex rational polyhedral cone.*

*Proof.* The maps $\partial$ and $H$ are defined over $\mathbb{Z}$, so $W$ is a rational linear subspace of $\mathbb{R}[E]$. The intersection of a rational linear subspace with the positive orthant is a convex rational polyhedral cone. □

**Lemma 3.5.** *Suppose that there exists a vector $w \in W$ with every component positive. Then $W^+$ has full rank in $W$.*

*Proof.* The condition implies that $W$ intersects the interior of $\mathbb{R}^+[E]$, which implies the result. □

**Definition 3.6.** A traintrack is *full* if it satisfies the hypothesis of Lemma 3.5. A traintrack is *recurrent* if there is a directed path from every vertex to every other vertex.

**Lemma 3.7.** *A traintrack is full if and only if each connected component is recurrent.*

*Proof.* The proof is obvious. □

An oriented immersion $S^1 \to T$ determines a weight in $\mathbb{Z}^+[E]$ whose coefficient for an edge $e$ is the degree with which $S^1$ maps over $e$; i.e. the degree of the map from $S^1$ to the oriented circle $T/(T-e)$.

**Definition 3.8.** We denote by $\mathcal{C}(T)$ the space of $\mathbb{R}$ linear formal sums of free homotopy classes of oriented immersions $S^1 \to T$ modulo the relation that if $\pi : L' \to L$ is an oriented cover of oriented 1-manifolds of degree $n$, then the composition $L' \to L \to T$ represents $n$ times $L \to X$.

We similarly define $\mathcal{C}^+(T)$ to be the positive orthant. Note that it does *not* make sense to define $\mathcal{C}(T)/\epsilon$. Composing $S^1 \to T$ with $h$ gives a map $S^1 \to X$, and we therefore obtain a linear map $h : \mathcal{C}(T) \to \mathcal{C}$ restricting to $h : \mathcal{C}^+(T) \to \mathcal{C}^+$. This map is typically neither injective nor surjective.

**Lemma 3.9.** *A weight $w \in \mathbb{Z}^+[E]$ is in the image of $\mathcal{C}^+(T)$ if and only if it is in $\ker \partial$.*

*Proof.* Represent $w$ as a finite collection of oriented edges. The condition $\partial w = 0$ is equivalent to the existence of a pairing of the vertices of this collection of edges, where each source vertex is paired with a target vertex and conversely. Gluing edges according to this pairing determines an oriented 1-manifold together with a tautological oriented immersion to $T$. The converse is obvious. □

**Lemma 3.10.** *A weight $w \in \mathbb{Z}^+[E] \cap \ker \partial$ is in $\ker H$ if and only if for any $c \in \mathcal{C}^+(T)$ representing $w$, the image of $c$ in $\mathcal{C}^+$ is null-homologous.*

*Proof.* This follows directly from the definition □

In words: an integral vector $w$ is in $W^+$ if and only if it is in the image of an integral $c \in \mathcal{C}^+(T)$ whose image in $\mathcal{C}^+$ is null-homologous.

3.2. **Arcs, rectangles and triangles.** We suppose given a traintrack $T$ with edges $E$, vertices $V$ and evaluation function $h$ as in § 3.1.

**Definition 3.11.** An *arc* is an ordered pair of vertices. If $\alpha$ is an arc, let $\overline{\alpha}$ denote the arc obtained by exchanging the order of the vertices (i.e. by reversing the "orientation" of $\alpha$). If $\mathbb{R}[A]$ denotes the $\mathbb{R}$ vector space spanned by arcs, let $\Lambda[A]$ denote the quotient by the relation $\overline{\alpha} = -\alpha$.

**Definition 3.12.** A *rectangle* is a cyclic list $(\alpha_0, e_0, \alpha_1, e_1)$ where the $\alpha_i$ are arcs and the $e_i$ are edges, for which $he_0 = \iota he_1$, and such that $t\alpha_i = se_i$ and $s\alpha_i = te_{i-1}$, indices taken mod 2.

**Definition 3.13.** A *triangle* is a cyclic list of arcs $(\alpha_0, \alpha_1, \alpha_2)$ such that $t\alpha_i = s\alpha_{i+1}$, indices taken mod 3.

**Definition 3.14.** Let $P$ denote the set of polygons (i.e. rectangles or triangles as above). There are linear maps $D_A : \mathbb{R}[P] \to \Lambda[A]$ and $D_E : \mathbb{R}[P] \to \mathbb{R}[E]$ defined on generators in the following way.

If $r = (\alpha_0, e_0, \alpha_1, e_1)$ is a rectangle, $D_A r = \alpha_0 + \alpha_1$ and $D_E r = e_0 + e_1$. If $\tau = (\alpha_0, \alpha_1, \alpha_2)$ is a triangle, $D_A \tau = \alpha_0 + \alpha_1 + \alpha_2$ and $D_E \tau = 0$.

We also use the notation $D := D_A \oplus D_E : \mathbb{R}[P] \to \Lambda[A] \oplus \mathbb{R}[E]$.

If $\tau = (\alpha_0, \alpha_1, \alpha_2)$ is a triangle, we define $\overline{\tau} := (\overline{\alpha}_2, \overline{\alpha}_1, \overline{\alpha}_0)$. Note that $D(\overline{\tau}) = -D(\tau)$.

**Definition 3.15.** Define the *polygon weight space* $Y$ to be the kernel

$$\widehat{Y} := \ker D_A : \mathbb{R}[P] \to \Lambda[A]$$

We define the *cone of positive weights* $Y^+ := \widehat{Y} \cap \mathbb{R}^+[P]$. Furthermore, define $Y \subset \mathbb{R}[P]$ to be the linear span of $Y^+$.

*Remark* 3.16. We do not assume that $Y^+$ has full rank in $\widehat{Y}$.

Let $\Sigma$ be an admissible surface, and suppose we have a map $\Sigma \to X$. Suppose further that $\partial\Sigma$ contains some components $\partial^i\Sigma$ (called *ideal components*) mapping to the basepoint of $X$, and that the restriction of the map to the other components $\partial^e\Sigma := \partial\Sigma - \partial^i\Sigma$ factors through an oriented immersion $\partial^e\Sigma \to T$. Choose such an immersion to $T$.

It is convenient to use the language of fatgraphs.

**Definition 3.17.** A *fatgraph* is a graph together with a choice of cyclic order of the edges incident to each vertex.

A fatgraph $Y$ can be canonically *fattened* to an oriented compact surface $\Sigma(Y)$ with boundary in which it sits as a *spine*; that is, in such a way that $\Sigma(Y)$ deformation retracts to $Y$. See Figure 1 for an example. For an introduction to fatgraphs, see e.g. [30] § 1. We will return to fatgraphs again in the sequel, especially in § 5.3 and § 7.4.



FIGURE 1. A fatgraph and its fattening

Now, suppose that $\Sigma$ as above is not compressible. Then $\Sigma$ has a *fatgraph representation*, i.e. there is some fatgraph $Y$ so that $\Sigma = \Sigma(Y)$, and a simplicial map from $Y$ to $X$ for which the given map from $\Sigma$ to $X$ factors through the deformation retraction to $Y$. The proof of this fact is implicit in Culler [17]; see also [8], especially § 4.1.

Each edge of the fatgraph determines a rectangle, and each vertex of the fatgraph determines a polygon which can be decomposed (efficiently) into triangles. Call the result a *triangulated fatgraph over* $T$.

**Definition 3.18.** We denote by $\mathcal{S}(T)$ the space of $\mathbb{R}$ linear formal sums of triangulated fatgraphs over $T$ modulo the relation that if $\pi : \Sigma' \to \Sigma$ is an oriented cover of degree $n$ (compatible with the triangulation and fatgraph structure) then $\Sigma'$ represents $n$ times $\Sigma$.

We similarly define $\mathcal{S}^+(T)$. There are tautological maps $\mathcal{S}(T) \to \mathcal{S}$ and $\mathcal{S}^+(T) \to \mathcal{S}^+$ given by forgetting the fatgraph structure and the choice of oriented immersion $\partial^e\Sigma \to T$. Note that the function $\chi$ on $\mathcal{S}^+$ pulls back to a well-defined function $\chi$ on $\mathcal{S}^+(T)$.

Decomposing a triangulated fatgraph into rectangles and polygons determines a map $\mathcal{S}(T) \to \mathbb{R}[P]$.

**Lemma 3.19.** *A weight $y \in 2\mathbb{Z}^+[P]$ is in the image of $\mathcal{S}^+(T)$ if and only if it is in $\ker D_A$. In particular, the image of $\mathcal{S}^+(T)$ in $\mathbb{R}[P]$ is exactly $Y^+$.*

*Proof.* One direction is obvious, since every weight in the image of $\mathcal{S}^+(T)$ is in $\ker D_A$. Conversely, let $y$ be even positive integral in $\ker D_A$. We consider a collection of rectangles and triangles determined by $y$, and try to glue boundary arcs in pairs.

By the definition of the vector space $\Lambda[A]$ and the fact that $y$ is even, the arcs can be paired and the polygons glued up to make a surface. Each rectangle admits a tautological map to $X$ coming from the edge labels, in such a way that the boundary arcs map to the basepoint. Map each triangle to $X$ by the constant map. The result is a compact oriented surface $\Sigma$ mapping to $X$. The components $\partial^e\Sigma$ admit a canonical lift to $T$ determined by the collection of edges. This completes the proof. $\square$

Given $\Sigma \to X$ representing a class in $\mathcal{S}^+(T)$, the boundary $\partial^e\Sigma \to T$ represents a class in $\mathcal{C}^+(T)$. This defines a linear map $\partial : \mathcal{S}^+(T) \to \mathcal{C}^+(T)$ compatible with $\partial : \mathcal{S}^+ \to \mathcal{C}^+$.

**Lemma 3.20.** *The image of $Y^+$ under $D_E$ is $W^+$. Moreover, if $T$ is full, then the image of $Y$ under $D_E$ is $W$.*

*Proof.* Every $y \in Y^+$ is in the image of $s \in \mathcal{S}^+(T)$. Evidently $D_E y$ is equal to the image of $\partial s \in \mathcal{C}^+(T)$ and is therefore in $W^+$ by Lemma 3.10. This proves one direction.

Conversely, every rational $w \in W^+$ is in the image of some $c' \in \mathcal{C}^+(T)$ whose image $c \in \mathcal{C}^+$ is in $B^+$. Therefore there is $s \in \mathcal{S}^+$ with $\partial s = c$ and after compressing $s$ if necessary, we can represent $s$ by a triangulated fatgraph. Because $\partial s$ is in the image of $\mathcal{C}^+(T)$, this triangulated fatgraph determines $s' \in \mathcal{S}^+(T)$ with $\partial s' = c'$, and therefore if $y \in Y^+$ is the image of $s'$ we have $D_E y = w$. This proves the first claim.

To prove the second claim, observe that if $T$ is full, then every $w \in W$ is of the form $w^+ - w^-$ for $w^\pm \in W^+$. Choose $y^\pm \in Y^+$ mapping to $w^\pm$ under $D_E$. Then $D_E(y^+ - y^-) = w$. $\square$

### 3.3. Euler characteristic.

**Definition 3.21.** Define $\chi : \mathbb{R}^+[P] \to \mathbb{R}$ to be 0 on rectangles, and $-1/2$ on triangles, and extend by linearity.

**Lemma 3.22.** *Suppose $y \in Y^+$ and let $s \in \mathcal{S}^+(T)$ map to $y$. Then $\chi(s) = \chi(y)$.*

*Proof.* By linearity, it suffices to prove this on generators. So let $\Sigma$ be a triangulated fatgraph over $T$. This fatgraph deformation retracts onto a spine, composed of an edge for each rectangle, and a tripod for each triangle. Edges and tripods are glued along 2-valent vertices (corresponding to paired arcs in $S$). Consequently the resulting graph is trivalent, with one vertex for each triangle. For a trivalent graph with $v$ vertices and $e$ edges, $e = 3v/2$ and $\chi = v - e = v - 3v/2 = -v/2$. ☐

Putting this together with our earlier observation that $\chi : \mathcal{S}^+(T) \to \mathbb{R}$ factors through $\mathcal{S}^+(T) \to \mathcal{S}^+$, we obtain a commutative diagram

$$\begin{tikzcd}
\mathcal{S}^+(T) \arrow[r] \arrow[d] & \mathcal{S}^+ \arrow[d, "\chi"'] \\
Y^+ \arrow[r, "\chi"] & \mathbb{R}
\end{tikzcd}$$

3.4. **Linear programming and rationality.** For $w \in W^+$, let $\mathcal{C}^+(w) \subset \mathcal{C}^+(T)$ denote its preimage in $\mathcal{C}^+(T)$. This is a convex set. Since the image of $\mathcal{C}^+(w)$ is contained in $B^+$, the function scl can be pulled back to $\mathcal{C}^+(w)$.

**Definition 3.23.** For $w \in W^+$, define $\mathrm{scl}(w)$ to be the infimum of $\mathrm{scl}(c)$ over all $c \in \mathcal{C}^+(w)$.

**Theorem 3.24** (Traintrack Rationality)**.** *For $w \in W^+$, there is a formula*

$$\mathrm{scl}(w) = \inf\{-\chi(y)/2 \ : \ y \in Y^+ \text{ and } D_E(y) = w\}$$

*Moreover, the infimum is achieved, and if $w$ is rational, the infimum is achieved at a rational $y$. Furthermore, the function* $\mathrm{scl}(w)$ *varies piecewise rational linearly on $W^+$.*

*Proof.* Since $W^+$ and $Y^+$ are rational polyhedral cones, the infimum is achieved, and if $w$ is rational, it is achieved at a rational $y$.

The function $\chi$ on $\mathcal{S}^+(T)$ pushes forward to $\chi$ on $\mathcal{S}^+$, and also pushes forward to $\chi$ on $Y^+$. Since every surface can be compressed (thereby reducing $-\chi$) until it is represented by a fatgraph, we conclude that for any homologically trivial $c \in \mathcal{C}^+$,

$$\mathrm{scl}(c) = \inf\{-\chi(s)/2 \ : \ s \in \mathcal{S}^+(T) \text{ and } \partial s = c\}$$

Taking a further infimum over all $c \in \mathcal{C}^+(T)$ mapping to $w$ we obtain the desired result.

The last assertion is just the observation that the output of linear programming with fixed variables depends piecewise rational linearly on the constraints. ☐

Note that the function $-\chi/2$ on $Y^+$ extends in an obvious way to a pseudonorm $\|\cdot\|$ on $\mathbb{R}[P]$ taking the value 0 on rectangles, and 1/4 on triangles. With this definition, $\mathrm{scl}(w) = \inf \|y\|$ over all $y \in Y^+$ with $D_E(y) = w$.

## 4. Counting functions and posimorphisms

In this section we consider a canonical family of traintracks associated to a free group $F$ and a choice of free generating set, namely the (so-called) *verbal traintracks*. Dual to the weights on a verbal traintrack are the *counting functions*, and the antisymmetric functions are exactly the (homogeneous) *counting quasimorphisms* considered by Rhemtulla [32] and Brooks [5].

We show that the system of (normed) weight cones $W_\ell^+$ on all the verbal traintracks together faithfully recover the geometry of the (infinite dimensional) normed cone $B^+$. Duality (i.e. the Hahn-Banach theorem) applied to these finite dimensional cones shows that for each $b \in B^+$ there is a rational counting function $\phi$ which, thought of as a linear function on $B^+$, has a level set that is a supporting hyperplane for the scl ball at $b/\text{scl}(b)$.

### 4.1. **Formalism.**

Suppose we have vector spaces $Y, W$ and a surjective linear map $D : Y \to W$. A pseudonorm $\| \cdot \|$ on $Y$ determines a pseudonorm $\| \cdot \|$ on $W$ by

$$\|w\| = \inf_{y \in D^{-1}(w)} \|y\|$$

Suppose further that we have convex cones $Y^+, W^+$ in $Y, W$ of full dimension, so that $DY^+ = W^+$. There is a different pseudonorm $\| \cdot \|^+$ on $W^+$ defined by

$$\|w\|^+ = \inf_{y \in D^{-1}(w) \cap Y^+} \|y\|$$

Since $D^{-1}(w) \cap Y^+$ is a subset of $D^{-1}$, there is an inequality

$$\|w\|^+ \geq \|w\|$$

but equality will not hold in general. Given $w$, the restriction of $\| \cdot \|$ to the affine subspace $D^{-1}(w)$ is convex, so if the infimum is not realized on the convex subset $Y^+ \cap D^{-1}(w)$, the infimum over this subset is achieved on the boundary. Conversely, if $\|w\|^+$ is achieved by some $y$ in the interior of $Y^+$, then $\|w\|^+ = \|w\|$.

### 4.2. **Verbal traintrack.**

**Definition 4.1.** Fix some length $\ell$. Define the *verbal traintrack of width* $\ell$ (denoted $T_\ell$) to be the following directed graph.

1. The vertices are the reduced words in $F$ of length $\ell - 1$.
2. The edges are the reduced words in $F$ of length $\ell$.
3. For an edge $e$ define $se$ and $te$ to be the prefix and suffix of $e$ of length $\ell - 1$.
4. For an edge $e$ define $he$ to be the first letter of $e$.

Note that $T_\ell$ is full for any $\ell$. For the remainder of this section we fix a width $\ell$ and work in the associated traintrack $T_\ell$.

*Remark* 4.2. Defining $he$ to be the first letter of $e$ is just a convention; if we fix some $i \leq \ell$ and define $he$ to be the $i$th letter of $e$, we obtain isomorphic normed spaces and cones, and moving back and forth between different conventions is straightforward (if somewhat messy).

We define $\mathcal{E}$ to be the vector in $\mathbb{R}[E]$ with coefficient 1 on every edge, and $\mathcal{R}$ the vector in $\mathbb{R}[P]$ with coefficient 1 on every rectangle.

**Lemma 4.3.** *$\mathcal{E} \in W^+$ and $\mathcal{R} \in Y^+$, and $D_E \mathcal{R} = (|S| - 1)^{\ell - 1} \mathcal{E}$.*

*Proof.* Let $e$ be an edge, with label $he$. There is a bijection between the set of $e'$ with $he' = \iota he$ and the set of rectangles $r$ with $D_E r = e + e'$. By the definition of $T_\ell$, given $e$ the number of such $e'$ is $(|S| - 1)^{\ell - 1}$ where $S$ is the symmetric generating set for $F$. $\square$

We use the notation $\|\cdot\|$ and $\|\cdot\|^+$ from § 4.1 for the pseudonorms on $W$ and $W^+$ associated to the pseudonorm on $Y$ taking the value $1/4$ on triangles and $0$ on rectangles. With this notation, $\|w\|^+ = \mathrm{scl}(w)$ for $w \in W^+$ and therefore $\mathrm{scl}(w) \ge \|w\|$. Furthermore, $\mathrm{scl}(\mathcal{E}) = \|\mathcal{E}\| = 0$. If we need to stress the dependence of $\|\cdot\|$ and $\|\cdot\|^+$ on $\ell$, we write $\|w\|_\ell$ and $\|w\|_\ell^+$.

**Lemma 4.4** (formula for norm)**.** *For any $w \in W^+$ there is a formula $\|w\| = \inf_t \mathrm{scl}(w + t\mathcal{E})$ over all $t \ge 0$. This infimum is achieved for all $t \ge t(w)$. Moreover, $t(w)$ depends piecewise rational linearly on $w$.*

*Proof.* First of all,

$$\|w\| \le \|w + t\mathcal{E}\| + \|-t\mathcal{E}\| = \|w + t\mathcal{E}\| \le \mathrm{scl}(w + t\mathcal{E})$$

so one direction is proved.

Conversely, Let $y$ be a vector in $D_E^{-1}(w)$, and write $y = y^+ - y^-$ for $y^\pm \in Y^+$ where we assume $y^+$ and $y^-$ have disjoint support in $P$. Let $x$ be the part of $y^-$ supported in triangles, and let $\overline{x}$ be obtained by replacing each $\tau$ in $x$ with $\overline{\tau}$. Then $D(x + \overline{x}) = 0$ so we can write $y' = (y^+ + \overline{x}) - (y^- - x)$ with $\|y'\| = \|y\|$ and $D_E(y') = y$. We must be careful, since $y^+ + \overline{x}$ is *not* necessarily in $Y^+$, although it is in $\mathbb{R}^+[P]$.

Now, since $y^- - x$ consists entirely of rectangles, there is $z \in \mathbb{R}^+[P]$ so that $y^- - x + z = t\mathcal{R}$ for some positive $t$. Hence we can write $y'' = (y^+ + \overline{x} + z) - (y^- - x + z)$ where now $(y^+ + \overline{x} + z)$ and $(y^- - x + z)$ are in $Y^+$, and $\|y''\| = \|y'\| = \|y\|$. Since $D_E(y^+ + \overline{x} + z) = w + t\mathcal{E}$ we have $\|y\| = \|y''\| \ge \mathrm{scl}(w + t\mathcal{E})$ and we deduce $\|w\| = \inf_t \mathrm{scl}(w + t\mathcal{E})$.

By construction, scl is piecewise rational linear and convex, and therefore the derivative $\frac{d}{dt}\mathrm{scl}(w + t\mathcal{E})$ is locally constant with finitely many jumps, and is strictly increasing at those jumps. On the other hand, $0 \le \mathrm{scl}(w + t\mathcal{E}) \le \mathrm{scl}(w)$ for all $t \in [0, \infty)$, so $\mathrm{scl}(w + t\mathcal{E})$ is eventually constant and equal to its infimum. If $w$ is rational, the infimal $t$ with this property is rational. □

### 4.3. Canonical immersion.

Let $T_\ell$ denote the verbal traintrack of length $\ell$. There is a canonical map $\Phi_\ell : C \to \ker \partial \subset \mathbb{R}[E]$ defined on a cyclicly reduced word $g$ by reading the subwords of length $\ell$ of the bi-infinite power $\dot{g}$ with initial letter contained in a fixed fundamental domain. This map restricts to surjections $\Phi_\ell : B \to W$ and $\Phi_\ell : B^+ \to W^+$.

Note for any $b \in B^+$ we have inequalities

$$\|\Phi_\ell(b)\| \le \|\Phi_\ell(b)\|^+ \le \mathrm{scl}(b)$$

On the other hand, we have the following estimate.

**Lemma 4.5.** *Let $b \in B^+$, and suppose that every primitive word in the support of $b$ has length at most $\ell - 1$. Then $\|\Phi_\ell(b)\|^+ = \mathrm{scl}(b)$.*

*Proof.* The condition implies that the support of $\Phi_\ell(b)$ is an embedded 1-manifold in $T_\ell$, whose components correspond to the components in the support of $b$. Consequently $\Phi_\ell^{-1}\Phi_\ell(b) \cap B^+ = b$ and the lemma follows from Theorem 3.24. □

We make the following definition:

**Definition 4.6.** Define $\epsilon : \mathbb{R}[E] \to \mathbb{R}[E]$ on generators by $e \to -e^{-1}$.

This might appear to cause confusion with $\epsilon : B \to B$ defined earlier; however, no confusion should arise in practice because of the following lemma:

**Lemma 4.7.** *For any $b \in B$ we have $\epsilon\Phi_\ell b = \Phi_\ell \epsilon b$. Moreover, $\epsilon\mathcal{E}_\ell = -\mathcal{E}_\ell$.*

*Proof.* These facts are immediate from the definitions. □

### 4.4. **Posimorphisms.**

We briefly interrupt the discussion to recall some elements of the dual theory to scl, which is to say, the theory of *quasimorphisms*.

**Definition 4.8.** Let $G$ be a group. A homogeneous quasimorphism is a function $\phi : G \to \mathbb{R}$ for which there is a least non-negative real number $D(\phi)$ (called the *defect*) so that

1. for any $g \in G$ and any $n \in \mathbb{Z}$ there is equality $\phi(g^n) = n\phi(g)$; and
2. for any $g, h \in G$ there is an inequality $|\phi(gh) - \phi(g) - \phi(h)| \le D(\phi)$.

The $\mathbb{R}$ vector space of homogeneous quasimorphisms is denoted $Q(G)$, or just $Q$ if $G$ is understood.

A homogeneous quasimorphism is necessarily a class function; together with the homogeneity condition, we see that a homogeneous quasimorphism descends to an $\mathbb{R}$-linear function $\phi : B/\epsilon \to \mathbb{R}$. Any $\mathbb{R}$-linear function $\phi : B/\epsilon \to \mathbb{R}$ extends to $C/\epsilon \to \mathbb{R}$, and any two extensions differ by an element of $H^1(G)$. We can give a simple alternate characterization of homogeneous quasimorphisms which underscores its relationship to scl.

**Lemma 4.9.** *An $\mathbb{R}$-linear function $\phi : B/\epsilon \to \mathbb{R}$ is (the restriction of) a homogeneous quasimorphism with defect $\le D(\phi)$ if and only if for every admissible surface $S$ there is an inequality*

$$\phi(\partial S) \le -\chi^-(S) \cdot D(\phi)$$

*Equivalently, for any $b \in B/\epsilon$,*

$$\phi(b) \le \mathrm{scl}(b) \cdot 2D(\phi)$$

*Proof.* The equivalence of the two inequalities follows from Proposition 2.12. The first inequality is vacuous when $S$ is closed or a disk. If $S$ is an annulus, $\phi(\partial S) = 0$ follows from homogeneity, since $\phi(b^{-1}) = -\phi(b)$. In every other case, $S$ can be cut up into thrice-punctured spheres. If $S = S_1 \cup S_2$ glued up along $\alpha$ (with opposite orientations) then $\phi(\partial S_1 + \partial S_2) = \phi(\partial S)$ (because $\phi(\alpha^{-1}) = -\phi(\alpha)$ by homogeneity), and the lemma is proved. □

The second inequality in the statement of the lemma shows that a homogeneous quasimorphism is precisely an element of the dual of $B/\epsilon$ with its scl pseudonorm, and that $2D$ is the operator norm. This gives a remarkably short proof of the following well-known duality theorem:

**Theorem 4.10** (Generalized Bavard Duality; [3, 8])**.** *Let $G$ be any group. Then for any $b \in B/\epsilon$, there is equality*

$$\mathrm{scl}(b) = \sup_\phi \phi(b)/2D(\phi)$$

*where the supremum is taken over all homogeneous quasimorphisms $\phi$.*

On the other hand, working with $B^+$ lets us consider a more general class of dual functions. Motivated by Lemma 4.9, we make the following definition:

**Definition 4.11.** Let $G$ be a group. An $\mathbb{R}^+$-linear function $\phi : B^+ \to \mathbb{R}$ is a *posimorphism* if there is some least non-negative number $D(\phi)$ (called the defect) so that for any $b \in B^+$ there is an inequality

$$\phi(b) \le \mathrm{scl}(b) \cdot 2D(\phi)$$

We denote the space of posimorphisms by $P(G)$, or just $P$ if $G$ is understood. Note that $P$ is an $\mathbb{R}^+$-module (i.e. a convex cone), and is *not* in general a vector space. The Hahn-Banach theorem implies that for any $b \in B$ there is some posimorphism $\phi$ with $D(\phi) = 1$ and $\phi(b) = 2\,\mathrm{scl}(b)$; we call any such $\phi$ *extremal* for $b$. In the next section we will see that the theory of traintracks can be used to construct an explicit extremal posimorphism, dual to any $b \in B^+(F)$ for $F$ a free group.

*Remark* 4.12. The relationship between quasimorphisms and posimorphisms is similar to the relationship between harmonic and subharmonic functions, or between martingales and submartingales.

### 4.5. Counting functions.

In this subsection we introduce the class of *counting functions*. The subclass consisting of *antisymmetric* counting functions (discussed in § 4.6) are the most well-known class of quasimorphisms on free groups. These counting quasimorphisms were introduced by Rhemtulla [32], and further developed by Brooks [5]. Similar functions were introduced by Epstein-Fujiwara [19], although with a subtly different definition. The Brooks-Rhemtulla quasimorphisms are sometimes called "big counting quasimorphisms" whereas the Epstein-Fujiwara quasimorphisms called "small counting quasimorphisms" if it is necessary to distinguish them.

The following problem is central, and well-known:

**Question 4.13** (Counting Quasimorphism Question)**.** *Does every chain in a free group admit an extremal counting quasimorphism?*

See e.g. Grigorchuk [23] for a partial discussion. We show that by changing the problem slightly, we can give a positive answer, in the form of the Counting Duality theorem (below), which says that every element in a free group admits an extremal counting *posimorphism*. Our analysis further clarifies the difference between posimorphisms and quasimorphisms, and leads to a suggestive thermodynamic model of the potential difference (see Remark 4.25).

We begin with the definition of counting functions.

**Definition 4.14** (Counting function)**.** If $\sigma$ is a reduced (noncyclic) word in $F$, the *counting function* $C^\sigma(\cdot)$ counts the number of copies of $\sigma$ in a cyclically reduced word. Hence for $g$ a reduced cyclic word, $C^\sigma(g)$ counts the number of copies of $\sigma$ in $g$.

The function $C^\sigma$ extends by linearity to $C^\sigma : C \to \mathbb{R}$. We let $C^*_\ell$ denote the linear span of the space of counting functions on all words $\sigma$ of length $\ell$, and we let $C^*_{\mathrm{count}}$ denote the union over all $\ell$. There is a map from $\mathbb{R}[F]$ to $C^*_{\mathrm{count}}$ taking $\sigma$ to $C^\sigma$ and extending by linearity, but this map is *not* injective. For example, every cyclically reduced word is either a power of $a$, or else each string of consecutive $a$'s starts and ends with one of $b$ or $B$; hence $C^{ab} + C^{aB} = C^{ba} + C^{Ba}$. In other words, the representation of some $\phi \in C^*_\ell$ as an expression of the form $\sum t_\sigma C^\sigma$ is not unique.

The next lemma shows that any counting function has a representative in which all the words $\sigma$ in the support can be taken to have some fixed length (equal to the supremum of the lengths of the words appearing in any representative).

**Lemma 4.15.** *If $\ell \le \ell'$ then $C^*_\ell$ is a subspace of $C^*_{\ell'}$.*

*Proof.* For every reduced word $\sigma'$ of length $\ell'$ let $\text{pre}_\ell(\sigma')$ denote the prefix of length $\ell$. Then

$$C^\sigma \to (2|S|-1)^{\ell-\ell'} \sum_{\text{pre}_\ell(\sigma')=\sigma} C^{\sigma'}$$

extends to an identification of $C^*_\ell$ with a subspace of $C^*_{\ell'}$. □

It is natural to think that there should be a strong connection between counting functions and verbal traintracks. This is summarized in the following lemma.

**Lemma 4.16.** *$C^*_\ell$ is the dual of $\Phi_\ell C$ and $C^*_\ell/H^1$ is the dual of $\Phi_\ell B = W_\ell$.*

*Proof.* It is obvious from the definition that $C^*_\ell$ factors through $\Phi_\ell$. Conversely, any linear functional on $\Phi_\ell C = \ker \partial \subset \mathbb{R}[E]$ extends to a linear functional on $\mathbb{R}[E]$ which can be thought of as an element of $C^*_\ell$ in an obvious way. □

The next lemma illustrates the relationship between counting functions and posimorphisms.

**Lemma 4.17.** *A counting function $\phi \in C^*_\ell$ is a posimorphism (with some defect) if and only if $\phi(c+c^{-1}) \le 0$ for all $c \in C$.*

*Proof.* The condition is necessary, since $c + c^{-1} \in B$, and $\text{scl}(c+c^{-1}) = 0$ for any $c \in C$. Conversely, let $b \in B$ be an arbitrary rational chain, and let $\Sigma$ be an extremal fatgraph for $b$, so that $\partial\Sigma$ represents an oriented cover of $b$ of some degree. By linearity it will suffice to consider the case $\partial\Sigma = b$, and we must prove that $\phi(b) \le -\chi(\Sigma)\cdot 2D(\phi)$ for some constant $D(\phi)$ depending only on $\phi$. If some component of $\Sigma$ is an annulus, its boundary is of the form $c+c^{-1}$, so without loss of generality we can assume that no component of $\Sigma$ is an annulus, and every vertex $v$ of the fatgraph has valence$(v) \ge 3$.

We suppose that we an write $\phi = \sum t_i C^{\sigma_i}$ for some real numbers $t_i$, where $\sigma_i$ ranges over $F_\ell$. We let $e_j$, $e_j^{-1}$ denote the set of (oriented) labels appearing as edges of $\Sigma$; i.e. the collection of cyclic words $b$ can be cut up at finitely many points into the collection of ordinary words $e_j$, $e_j^{-1}$. It makes sense to evaluate a counting function on an ordinary (non-cyclic) word, by counting copies of subwords $\sigma_i$. The only copies of $\sigma_i$ that occur in $\partial\Sigma$ but not in some $e_j$ or $e_j^{-1}$ are those that map over some vertex of the fatgraph. Hence

$$|\phi(b) - \sum_j (\phi(e_j) + \phi(e_j^{-1}))| \le \sum_v \text{valence}(v)\cdot\ell\cdot\sum_i |t_i|$$

On the other hand, for each $j$ the difference of $\phi(e_j)$ as an ordinary word and as a cyclic word is also bounded by $\ell\cdot\sum_i |t_i|$, and therefore we can estimate

$$\phi(b) \le 2\sum_v \text{valence}(v)\cdot\ell\cdot\sum_i |t_i|$$

Since every vertex $v$ has valence at least 3, and since $-\chi^-(\Sigma) = \sum_v(\text{valence}(v)-2)/2$, it follows that we can estimate

$$\phi(b) \le -12\chi^-(\Sigma)\cdot\ell\cdot\sum_i |t_i|$$

so that $\phi$ is a posimorphism with $D(\phi) \le 12\ell \cdot \sum_i |t_i|$. □

*Remark* 4.18. We can give a simpler, although less self-contained or direct proof of Lemma 4.17 using the theory of verbal traintracks already developed. Evidently, $\phi(b) = \phi(\Phi_\ell(b))$ for any $b \in B$, so to show that $\phi$ is a posimorphism, it suffices to show that $\phi(w) \le \mathrm{scl}(w) \cdot 2D(\phi)$ for all $w \in W_\ell^+$ and for some $D(\phi)$. However, since $W_\ell^+$ is finite dimensional and $\phi$ is linear, it suffices to have $\phi(w) \le 0$ whenever $\mathrm{scl}(w) = 0$. Now, if $\mathrm{scl}(w) = 0$, there is some $b \in \Phi_\ell^{-1}(w)$ with $\mathrm{scl}(b)$, and every such $b$ is of the form $c + c^{-1}$. The proof follows.

Finally we come to the counting duality theorem, which explicitly constructs a rational extremal counting posimorphism dual to any $b \in B$.

**Theorem 4.19** (Counting Duality). *For any $b \in B^+$ there is an $\ell$ and a rational linear function $\phi \in C_\ell^*/H^1$ so that $\phi(b') \le \|\Phi_\ell(b')\|^+ \le \mathrm{scl}(b')$ for all $b' \in B^+$ and $\phi(b) = \mathrm{scl}(b)$. In other words, $\phi$ is an extremal posimorphism for $b$.*

*Proof.* We take $\ell$ larger than any word in the support of $b$, so that $\|\Phi_\ell(b)\|^+ = \mathrm{scl}(b)$ by Lemma 4.5. Consider the unit ball in the pseudo-norm $\|\cdot\|^+$ on $W_\ell^+$. This is a closed, convex, rational polyhedron. Consequently there is a rational supporting hyperplane which intersects the boundary of the unit ball along the ray $\mathbb{R}^+\Phi_\ell(b)$. Such a hyperplane is the level set $\phi = 1$ for some $\phi \in C_\ell^*$, by Lemma 4.16. By construction, $\phi(b') \le \|\Phi_\ell(b')\|^+ \le \mathrm{scl}(b')$ for all $b' \in B^+$, but $\phi(b) = \|\Phi_\ell(b)\|^+ = \mathrm{scl}(b)$. □

## 4.6. Antisymmetrization.

We now spell out the relationship between the norm $\|\cdot\|_\ell$ and the space of (homogeneous) counting *quasimorphisms*, which are precisely the antisymmetric counting functions.

**Definition 4.20.** A counting function $\phi$ is *antisymmetric* if $\phi(g^{-1}) = -\phi(g)$ for all $g$. An antisymmetric counting function is called a *counting quasimorphism*. The space of counting quasimorphisms is denoted $Q_{\mathrm{count}}$, and its intersection with $C_\ell^*$ is denoted $Q_\ell$.

**Lemma 4.21.** *For any $b \in B$ there is an equality $\|\Phi_\ell b\|_\ell = \frac{1}{2}\|\Phi_\ell(b + \epsilon b)\|_\ell$.*

*Proof.* For any $b \in B^+$ the chain $b - \epsilon b \in B^+$ is the union of two copies of $b$ with opposite orientations. Evidently there are equalities

$$0 = \mathrm{scl}(b - \epsilon b) = \|\Phi_\ell(b - \epsilon b)\|_\ell^+ = \|\Phi_\ell(b - \epsilon b)\|_\ell$$

for any $\ell$. By linearity, the same is true for any $b \in B$. We estimate

$$\begin{aligned} \|\Phi_\ell(b + \epsilon b)\|_\ell &= \|\Phi_\ell(b + \epsilon b)\|_\ell + \|\Phi_\ell(b - \epsilon b)\|_\ell \\ &\ge \|\Phi_\ell(b + \epsilon b + b - \epsilon b)\|_\ell = \|\Phi_\ell(2b)\|_\ell \\ &= \|\Phi_\ell(2b)\|_\ell + \|\Phi_\ell(-b + \epsilon b)\|_\ell \\ &\ge \|\Phi_\ell(2b - b + \epsilon b)\|_\ell = \|\Phi_\ell(b + \epsilon b)\|_\ell \end{aligned}$$

Each inequality above is therefore an equality, and the lemma is proved. □

It follows that $\|\Phi_\ell b\|_\ell$ is well-defined on $B/\epsilon$. This is significant, because $\Phi_\ell$ is *not* well-defined on $B/\epsilon$.

**Definition 4.22.** Define $\Psi_\ell : C \to \mathbb{R}[E]$ by $\Psi_\ell(c) = \frac{1}{2}\Phi_\ell(c + \epsilon c)$.

Note that $\|\Phi_\ell(b)\|_\ell = \|\Psi_\ell(b)\|_\ell$ for any $b \in B$, by Lemma 4.21.

**Lemma 4.23.** *$Q_\ell$ is the dual of $\Psi_\ell C$ and $Q_\ell/H^1$ is the dual of $\Psi_\ell B = (\mathrm{id}+\epsilon)W_\ell$.*

*Proof.* This follows just as in the proof of Lemma 4.16, together with the observation that the adjoint of $c \to \frac{1}{2}(c+\epsilon c)$ on $C$ is the "antisymmetrization map" $\epsilon^* : C^\sigma \to \frac{1}{2}(C^\sigma - C^{\sigma^{-1}})$ on $C^*_\ell$. □

The retraction $W_\ell \to \frac{1}{2}(\mathrm{id}+\epsilon)W_\ell$ factors through the quotient $W_\ell \to W_\ell/\epsilon$, and shows that this quotient has a section. We have shown that $\|\cdot\|_\ell$ descends to a pseudo-norm on $W_\ell/\epsilon$, and that the dual of $W_\ell/\epsilon$ is isomorphic to $Q_\ell/H^1$ by the adjoint of $\Psi_\ell$.

**Theorem 4.24** (Counting Quasimorphism Duality)**.** *For any $\ell$ and any $b \in B/\epsilon$, there is an equality*

$$\|\Psi_\ell(b)\|_\ell = \sup_{\phi \in Q_\ell/H^1} \frac{\phi(b)}{2D(\phi)}$$

*Proof.* The map $\Psi_\ell$ from $B/\epsilon$ to $W_\ell/\epsilon$ pulls back the pseudonorm $\|\cdot\|_\ell$ to a pseudonorm on $B/\epsilon$ which satisfies $\|\Psi_\ell(b)\|_\ell \le \mathrm{scl}(b)$. Moreover, for every $w \in W/\epsilon$ there is some $b' \in B/\epsilon$ with $\Psi_\ell(b') = w$ and $\|w\|_\ell = \mathrm{scl}(b')$.

Choose a supporting hyperplane for the unit ball in the pseudonorm $\|\cdot\|_\ell$ on $W/\epsilon$ that intersects the boundary of the ball at $\Psi_\ell(b)$. Let $b' \in B/\epsilon$ be such that $\Psi_\ell(b') = \Psi_\ell(b)$ and $\mathrm{scl}(b') = \|\Psi_\ell(b)\|_\ell$. Thinking of $Q_\ell/H^1$ as the dual of $W/\epsilon$, this supporting hyperplane is the level set of some $\phi \in Q_\ell/H^1$. We have $\phi(b') = \mathrm{scl}(b')$ and $|\phi(b'')| = \|\Psi_\ell(b'')\|_\ell \le \mathrm{scl}(b'')$ for every $b'' \in B/\epsilon$. Since $\mathrm{scl}(gh-g-h) \le 1/2$ for all $g,h$ we have $D(\phi) \le 1/2$. On the other hand, $|\phi(gh-g-h)| = 1/2$ whenever the thrice-punctured sphere bounded by $gh-g-h$ immerses in an extremal surface for $b'$. □

Compare with the Generalized Bavard Duality Theorem (i.e. Theorem 4.10). We deduce that a chain $b$ admits an extremal counting quasimorphism if and only if $\|b\|_\ell = \|b\|_\ell^+$ for some $\ell$.

*Remark* 4.25. For $b \in B/\epsilon$ we may suggestively refer to the quantity $\sup_\ell \|\Psi_\ell(b)\|_\ell$ as *renormalized* scl, in contrast to *bare* scl (i.e. to $\mathrm{scl}(b)$ itself). As explained in [8], § 4.3.6 there is a thermodynamic interpretation of scl, where labeled strands combine into configurations (i.e. fatgraphs) in such a way as to minimize energy (as measured by $-\chi$). If $b$-strands are inserted into a "Dirac sea" of free strands, the renormalized energy may be less, because fatgraphs may form that pair $b$ with collections of free strands that are themselves capable of being paired into parallel strands at increasing scales $\ell \to \infty$.

## 5. The geometry of $Q_\ell$

The theory of counting quasimorphisms as described in § 4 has the potential to build a bridge between topology and combinatorics. In this section we describe the geometry of $Q_\ell/H^1$ as a normed space with its defect norm, for free groups of finite rank, with special attention to the case of $\ell = 2$.

### 5.1. **Fatgraphs, ends, fat folding.**

In the sequel we assume the reader is familiar with Stallings' theory of graphs and graph foldings; see [36].

**Definition 5.1.** A *fatgraph with ends* is a graph with some 1-valent vertices (the ends). The fattening of a fatgraph with ends is a surface with corners; there is one edge of the boundary associated to each end of the graph; these edges are called *infinitesimal edges*.

**Definition 5.2.** Let $X$ be a graph. A *fatgraph over* $X$ is a fatgraph $Y$ together with a simplicial map $Y \to X$. A fatgraph over $X$ is *folded* if the induced map $\partial\Sigma(Y) \to X$ is an immersion.

The most important case to consider is when $X$ is a standard rose for a free group $F$ with a fixed (symmetric) generating set $S$. A fatgraph $Y$ over the standard rose is also called a *labeled fatgraph*. We can think of it as being given by assigning a reduced word to each oriented edge of $Y$. The fatgraph $Y$ is folded if the induced labels on $\partial\Sigma(Y)$ are cyclically reduced.

A simplicial map $Y \to X$ between graphs which is not an immersion (i.e. which is not locally injective) can be iteratively *Stallings folded*, by identifying pairs of oriented edges of $Y$ with the same initial vertex which map to the same oriented edge of $X$. If $Y$ is a fatgraph, a Stallings folding is a *fat folding* if the edges to be identified are adjacent in the cyclic order at the initial vertex. The result of a fat folding is a new fatgraph. A sequence of fat foldings is *elementary* if it induces a homotopy equivalence of fatgraphs. Every fatgraph $Y$ over a graph $X$ may be fat folded to produce a canonical folded fatgraph $Y'$ over $X$.

If $F$ is a free group with symmetric generating set $S$, and $X$ is a standard rose for $F$, a fatgraph $Y$ over $X$ is also called a *labeled fatgraph*. An oriented edge of $Y$ determines a word in $S$. If $Y$ is folded (as a fatgraph), the resulting labels on $\partial\Sigma(Y)$ are cyclically reduced. If $Y$ is a fatgraph with ends, $\partial\Sigma(Y)$ is labeled with reduced cyclic words, and reduced non-cyclic words starting and ending at an infinitesimal edge.

Now, suppose $Y', Y$ are two folded fatgraphs over $X$. We say that $Y'$ *immerses* in $Y$ *as a fatgraph* if there is a fatgraph $Y''$ that immerses in $Y$ in the usual sense (i.e. as an immersion of graphs, compatibly with the cyclic orders at each vertex), and $Y'$ is obtained by *elementary* fat folding $Y''$. Note that this implies that $Y'$ and $Y''$ are homotopic, and that the surface $\Sigma(Y'')$ which is homeomorphic to $\Sigma(Y')$ immerses in $\Sigma(Y)$ (in the usual sense). The point of this definition is that even though $Y$ might be folded over $X$ as a fatgraph, it might *not* be folded as an ordinary graph, and therefore there might be subfatgraphs which are not themselves folded.

The following theorem is a restatement of Theorem B from [10]:

**Theorem 5.3.** *Let $\Gamma$ be a chain in $B/\epsilon$, and let $X$ be a labeled fatgraph so that $\partial\Sigma(X) = \Gamma$, and $\Sigma(X)$ is an extremal surface for $\Gamma$. Let $X'$ be any fatgraph that immerses in $X$, and let $\Gamma' = \partial\Sigma(X')$. Then $\Sigma(X')$ is an extremal surface for $\Gamma'$. Moreover, if $\phi$ is any quasimorphism extremal for $\Gamma$, then $\phi$ is extremal for $\Gamma'$.*

### 5.2. **Tripods.**

A *tripod* $T$ is a fatgraph with three ends; its underlying graph is the cone on three (cyclically ordered) points. We consider tripods which are folded over the standard rose. Hence if $x, y, z$ are the three labels on the edges of the tripod (in positive cyclic order) then $\partial\Sigma(T) = Xy + Yz + Zx$. A tripod has *length* $\ell - 1$ if the words $x, y, z$ all have length $\ell - 1$.

Now, suppose $\phi \in Q_\ell$ is a counting quasimorphism, and let $T_{x,y,z}$ be a tripod of length $\ell - 1$. We define

$$\phi(T_{x,y,z}) := \phi(\partial\Sigma(T_{x,y,z})) = \phi(Xy + Yz + Zx)$$

**Lemma 5.4.** *Let $\phi \in Q_\ell$. Then $D(\phi) = \max_{x,y,z} 2\phi(T_{x,y,z})$ where the supremum is taken over all tripods of length $\ell - 1$.*

*Proof.* By counting duality (i.e. Theorem 4.24) there is some $\Gamma$ with $\phi(\Gamma)/2D(\phi) = \text{scl}(\Gamma)$. After replacing $\Gamma$ by a finite cover if necessary, there is a labeled fatgraph $X$ such that $\partial\Sigma(X) = \Gamma$, and $\text{scl}(\Gamma) = -\chi(X)/2$. After replacing $\phi$ by a multiple if necessary, we can assume $D(\phi) = 1$, so that $\phi(\Gamma) = -\chi(X)$. Now, let $T$ be any tripod of length $\ell - 1$ that immerses in $X$. We can build a new fatgraph $Y$ homotopic to a pair of pants that immerses in $X$, from two copies of $T$ joined by three edges of length $\geq 2\ell$. By Theorem 5.3 and by construction,

$$1/2 = -\chi(Y)/2 = \text{scl}(\partial\Sigma(Y)) = \phi(\partial\Sigma(Y))/2 = \phi(T)$$

This proves that $D(\phi) \geq \max_T 2\phi(T)$. Conversely, if $T$ is any tripod of length $\ell - 1$, we can build a fatgraph $Y$ from two copies of $T$ joined by edges of length $\geq 2\ell$. Then

$$\phi(T) = \phi(\partial\Sigma(Y))/2 \leq \text{scl}(\partial\Sigma(Y)) \leq -\chi(Y)/2 = 1/2$$

□

We can therefore define for each $T$ of length $\ell - 1$ the *tripod hyperplane* $\widehat{\pi}_T$ to be the affine subspace of $Q_\ell$ consisting of $\phi$ with $\phi(T) = 1/2$. By the definition of defect, and Lemma 5.4 we deduce the following:

**Lemma 5.5.** *The tripod hyperplanes $\widehat{\pi}_T$ as $T$ ranges over the tripods of length $\ell - 1$ are a complete set of supporting hyperplanes for the unit ball in $Q_\ell/H^1$.*

For each tripod $T$ of length $\ell - 1$, define the *tripod face* $\pi_T$ to be the intersection of $\widehat{\pi}_T$ with the boundary of the unit ball in $Q_\ell/H^1$.

It follows in particular that every codimension one face of the unit ball is a tripod face. However, it is by no means clear which tripods correspond to codimension one faces of the unit ball. In the next few sections we will show that every $\pi_T$ is nonempty, and in the special case of $\ell = 2$ that every $\pi_T$ is a codimension 1 face.

**Corollary 5.6.** *Let $\Gamma \in B/\epsilon$ be arbitrary, and let $X$ be a fatgraph whose fattening is an extremal surface for $\Gamma$. Fix some $\ell$, let $T_i$ denote the set of tripods of length $\ell - 1$ that immerse in $X$, let $\widehat{\pi}_i$ be the associated tripod hyperplanes, and let $\pi_i$ be the tripod faces. The set of $\phi \in Q_\ell/H^1$ extremal for $\Gamma$ is precisely equal to the cone on $\cap_i \pi_i$.*

*Proof.* This follows immediately from Theorem 5.3 and Lemma 5.5, modulo the following claim. We claim that we can find a collection $\{i_j\}$ of $-2\chi(X)$ tripods $T_{i_j}$ immersing in $X$ so that $\Psi_\ell(\Gamma) = \sum_j \Psi_\ell(T_{i_j})$ (see Definition 4.22 for the definition of $\Psi_\ell$). Assuming this claim, the converse is proved as follows. Any $\phi \in \cap_i \pi_i$ satisfies $D(\phi) = 1$ and $\phi(T_i) = 1/2$ for every tripod immersing in $X$. In particular, if such a $\phi$ exists, then $\phi(\Gamma)/2D(\phi) = \sum_j \phi(T_{i_j})/2 = -\chi(X)/2 = \text{scl}(\Gamma)$ so $\phi$ is extremal for $\Gamma$.

The claim is proved by a standard argument in hyperbolic geometry. Pick a hyperbolic metric on $\Sigma(X)$ with geodesic boundary, and fix an ideal triangulation of this surface in which the ideal triangles spin around the boundary. Now deform the metric so that the length function (projectively) degenerates to the length function on the graph $X$. After rescaling the metric, the diameter of the thick part of each ideal triangle goes uniformly to zero, and the center of this thick part converges to one of the vertices of $X$. Cut off the ends of each triangle so that the length of each end is equal to $\ell - 1$ in the rescaled metric. Each triangle degenerates to a tripod. This collection of tripods evidently satisfies the claim. $\square$

We call a collection $T_i$ of tripods of length $\ell - 1$ *compatible* if the associated tripod faces $\pi_i$ have a nonempty intersection in the boundary of the unit ball in $Q_\ell/H^1$. Corollary 5.6 implies that any fatgraph built from copies of a compatible collection of tripods will be extremal for its boundary.

### 5.3. Basic realizations and rotation quasimorphisms.

Fix a free group $F$ of finite rank, and a finite symmetric generating set $S$. If the rank of $F$ is $k$, then $|S| = 2k$. We fix a standard rose $X$, i.e. a graph with one vertex $*$ and $k$ edges, and an identification of $F$ with $\pi_1(X, *)$ in such a way that the elements of $S$ correspond to the oriented edges of $X$.

A cyclic ordering $O$ on $S$ gives $X$ the structure of a fatgraph with fattening $\Sigma_O$ (see Definition 3.17) and thereby determines an isomorphism from $F$ to $\pi_1(\Sigma_O)$ up to conjugacy. We call such an isomorphism a *basic realization* with basis $S$.

The embedding of $X$ in $\Sigma$ determines an embedding of the universal cover $\widetilde{X}$ in $\widetilde{\Sigma}$, which is homeomorphic to the (oriented) plane, and this embedding determines a cyclic ordering on $\mathcal{E}(\widetilde{X})$, the space of *ends* of $\widetilde{X}$. Since $H^2(F) = 0$, this action on a cyclically ordered set determines a *rotation quasimorphism* $\text{rot}_O \in Q/H^1$; see [8] § 2.3.3 for details.

In fact, for the rotation quasimorphisms arising from basic realizations as above, it turns out that $\text{rot}_O \in Q_2/H^1$, and we can give an explicit formula for a representative. Recall that we abbreviate inverse of a generator (given as a lower case letter) by capitalization, so that $X$ means $x^{-1}$, and so on.

**Theorem 5.7** (Rotation number formula)**.** *Let $O$ be a cyclic order on $S$. For each pair of letters $x, y \in S$ with $y$ not equal to $X$, there is a* linear order $O|S - X$ *on $S - X$, and we let $n_{xy}$ denote the (signed) distance from $x$ to $y$ in this linear order. Then there is a formula*

$$\text{rot}_O = \frac{1}{2k} \sum_{y \neq X} n_{xy} C^{xy}$$

See [8] Thm. 4.76 for a proof.

*Remark* 5.8. Some version of this formula was originally discovered by Chillingworth [15], although it was expressed in quite different language, and he did not make the connection to the theory of quasimorphisms.

In [10] it is proved that rotation quasimorphisms associated to realizations are extremal vertices in the space of all homogeneous quasimorphisms; i.e. that they are dual to codimension one faces of the scl norm on $B/\epsilon$. In particular it follows that they determine vertices of the unit ball in $Q_\ell/H^1$ for any $\ell \geq 2$.

We now relate rotation quasimorphisms to tripod faces.

**Lemma 5.9.** *Let $T$ be a tripod face of length $\ell - 1$, and $\text{rot}_O$ a rotation quasimorphism. Let $x, y, z$ be the initial letters of the edge labels on $T$. Then $\pi_T$ contains $\text{rot}_O$ if and only if $x, y, z$ are positively circularly ordered in $O$.*

*Proof.* This follows immediately from the definition, and from the formula in Theorem 5.7. Alternately, if $Y$ is a fatgraph made from two copies of $T$, then either $Y$ or $Y$ with the opposite orientation immerses in $X$ with the fatgraph structure associated to $O$. The surface $\Sigma(X)$ is extremal for $\partial\Sigma(X)$, and $\text{rot}_O$ is extremal for $\partial\Sigma(X)$. Now apply Theorem 5.3. $\square$

5.4. **The geometry of $Q_2$.** Let $\Delta$ denote the complete simplex on $S$, a symmetric free generating set for $F$. Let $\widetilde{C}^*$ denote the complex of simplicial (real-valued) cochains on $\Delta$. For each $i$, the symmetric group $S_{i+1}$ acts on $\widetilde{C}^i$ by permuting the vertices of an $i$-simplex; call a cochain *antisymmetric* if it transforms by the sign character under the action of $S_{i+1}$. The subcomplex $C^*$ of antisymmetric cochains includes into $\widetilde{C}^*$, and this inclusion is a homotopy equivalence (over $\mathbb{Q}$). Note that every element of $C^*$ vanishes on degenerate simplices.

Since $\Delta$ is contractible, we have $Z^i = B^i$ for all $i > 0$; i.e. every cocycle is a coboundary.

**Lemma 5.10.** *There is an isomorphism of vector spaces $Z^2 = Q_2/H^1$. Moreover, the defect norm on $Q_2/H^1$ is twice the $L^\infty$ norm on $Z^2$.*

*Proof.* First of all, note that an element of $Q_2/H^1$ is determined by its values on tripods of length 1. This follows from Lemma 5.5 since the tripod hyperplanes are a complete set of supporting hyperplanes, and since the defect norm on $Q_2/H^1$ is nondegenerate.

An ordered pair of elements $s, t \in S$ determines a word $s^{-1}t$ of length 2 in $F$. Insisting that $s, t$ are distinct is equivalent to insisting that $s^{-1}t$ is reduced. Therefore an element of $Q_2$ (i.e. an antisymmetric counting function of length 2) is exactly the same thing as an antisymmetric simplicial 1-cochain, i.e. an element of $C^1$. If $\phi$ is a 1-cochain, then $\delta\phi(s,t,u) = \phi(s,t) + \phi(t,u) + \phi(u,s)$. If we think of $s, t, u$ as the ordered labels on a tripod $T$ of length 1, then $\delta\phi(s,t,u) = \phi(\partial\Sigma(T))$ where on the right hand side we think of $\phi$ as a counting quasimorphism. Thus the image of $\phi$ in $Q_2/H^1$ is precisely determined by its image in $Z^2$, and since $B^2 = Z^2$, every element of $Q_2/H^1$ arises in this way. Finally, equating defect with twice the $L^\infty$ norm is just a restatement of Lemma 5.4 in this context. $\square$

From this we reduce a strong structure theorem for the unit ball in $Q_2/H^1$:

**Theorem 5.11** (Unit Ball)**.** *The codimension 1 faces of the unit ball in $Q_2/H^1$ are exactly the set of length 1 tripod faces. The symmetric group $S_{2k}$ acts on this set by permutation of edge labels, and the resulting action on $Q_2/H^1$ is by isometries. As a representation of $S_{2k}$, the space $Q_2/H^1$ is isomorphic to $\wedge^2 V$ where $V$ is the standard $(2k-1)$-dimensional representation.*

*Proof.* The group $S_{2k}$ acts on $\Delta$ by permutations, and therefore on $Q_2/H^1$ by isometries, by Lemma 5.10. Now, if $\widehat{V}$ denotes the (coordinate) permutation representation of $S_{2k}$ on $\mathbb{R}^{2k}$, there is an obvious identification of $C^0$ with $\widehat{V}$ and $C^1$ with $\wedge^2\widehat{V}$. Then $Z^2 = C^1/\delta C^0 = \wedge^2 V$ where $V$ is the complement in $\widehat{V}$ of the constant vector, which is the kernel of $\delta : C^0 \to C^1$. This shows that $Q_2/H^1$ is isomorphic to $\wedge^2 V$ as a vector space, and that the action of $S_{2k}$ is irreducible. It follows that the rotation quasimorphisms $\text{rot}_O$ span $Q_2/H^1$. Since every tripod face $\pi_T$ contains exactly half of the $\text{rot}_O$ by Lemma 5.9, it follows that every tripod face is a codimension one face. $\square$

*Remark* 5.12. The group $\text{Out}(F)$ acts by isometries on $Q/H^1$, and the stabilizer of a symmetric generating set $S$ therefore acts by isometries on $Q_2/H^1$. This stabilizer is the group $\mathbb{Z}/2\mathbb{Z}^k \rtimes S_k$, where $S_k$ acts on $F$ by permuting the generators, and the

$\mathbb{Z}/2\mathbb{Z}$ factors act by switching a generator with its inverse. Note that this group acts reducibly on $Q_2/H^1$ in the case $F$ has rank 2. It is therefore quite surprising that $Q_2/H^1$ admits the full group $S_{2k}$ as isometries; this seems quite mysterious from the topological perspective, since such permutations do not even preserve the topological type of a fattening!

*Remark* 5.13. Since $Q_2/H^1$ includes as a subspace of $Q_\ell/H^1$ for each $\ell \geq 2$, and since the rotation quasimorphisms are vertices in each $Q_\ell/H^1$, it follows that every tripod face $\pi_T$ is nonempty, and in fact has dimension at least $2k^2 - 3k$. On the other hand, for $\ell \geq 3$ it is *not* true that every tripod face has codimension one. See Example 5.19.

## 5.5. **Examples.**

*Example* 5.14 ($Q_2$ in rank 2). For $k = 2$ the dimension of $Q_2/H^1$ is 3. The unit ball is an octahedron, whose extremal vertices correspond to the six distinct cyclic orders on the set of generators and their inverses; see Figure 2. These correspond (in antipodal pairs, one for each orientation) to three realizations of $F_2$ as $\pi_1(\Sigma)$: one as the once-punctured torus with boundary representing the commutator $[x, y]$, and two as thrice-punctured spheres, in the first case with boundary representing the conjugacy classes $x, y, XY$ and in the second case with boundary representing the conjugacy classes $x, Y, Xy$.



FIGURE 2. The unit ball in $Q_2/H^1$ for rank 2.

*Example* 5.15 ($Q_2$ in rank 3). For $k = 3$ the dimension of $Q_2/H^1$ is 10. The unit ball has 120 vertices and 40 faces, which are precisely the set of tripod faces. Tripod faces come in antipodal pairs; each face contains exactly 60 vertices, so an antipodal pair contain all the vertices. The vertices are exactly the set of rotation quasimorphisms associated to basic realizations.

*Example* 5.16 ($Q_2$ in rank 4). For $k = 4$ the dimension of $Q_2/H^1$ is 21. There are 112 tripod faces, which come in 56 antipodal pairs. There are 267120 vertices, of which 5040 are the rotation vertices.

What about the 262080 non-rotation vertices in rank 4? Notice that every rotation vertex takes the value $1/2$ or $-1/2$ on every tripod. It turns out experimentally that every vertex of the unit ball in $Q_2/H^1$ in rank 4 takes values in $\{-1/2, 0, 1/2\}$ on every tripod, and in fact this holds for all ranks.

**Proposition 5.17.** *For any rank, every vertex of the unit ball in $Q_2/H^1$ takes values in $\{-1/2, 0, 1/2\}$ on every tripod.*

*Proof.* This follows from general facts about 2-cocycles on contractible simplicial complexes. Recall Lemma 5.10 equating $Q_2/H^1$ and $Z^2$ of the simplex $S$ whose vertices are the symmetric generating set of the free group. Let us be given a vertex of the unit ball $\phi \in Q_2/H^1$. By abuse of notation, we'll also denote the function on 1-simplices by $\phi$. Because $\phi$ is a vertex, it is determined by the collection of tripods on which it takes $1/2$ its defect (which must be 1). Equivalently, $\delta\phi$ is determined by the 2-simplices on which it is $1/2$. We will try to conclude the value of $\phi$ from the known values of $\delta\phi$. Consider any connected collection $K$ of simplices on which the value of $\delta\phi$ is fixed (at $1/2$). We can set the value of $\phi$ on each 1-simplex in $K$ to be 0 or a sum or difference of values of $\delta\phi$, as appropriate to get the correct values of $\delta\phi$ on $K$. The fact that $\delta\phi$ is a cocycle ensures that this is possible. Now, these values for $\phi$ may not be extendable over the entire simplex $S$, but they extend *up to a* 1*-cocycle on* $K$.

Now take any 2-simplex $s$ on which the value of $\delta\phi$ is not $\pm 1/2$; to prove the proposition, we must show that $\delta\phi(s) = 0$. Note that $|\delta\phi(s)| < 1/2$, because by assumption $\phi$, being a vertex, takes its defect on a maximal collection of tripods. Take a collection $K$ of 2-simplices on which $\delta\phi$ is $\pm 1/2$, such that this collection includes the 1-simplices in $s$. This is possible because the values of $\phi$ must be determined by $\delta\phi$. As above, we can assign values to $\phi$ compatibly, which will be correct up to a 1-cocycle on $K$, and these values will be sums and differences of the values of $\delta\phi$; that is, they will be in $\frac{1}{2}\mathbb{Z}$. We can compute $\delta\phi(s)$ using these values of $\phi$, so $\delta\phi(s) \in \frac{1}{2}\mathbb{Z}$, so we have $\delta\phi(s) = 0$, as desired. $\square$

A similar phenomenon occurs in rank 2 for $\ell = 3$, and we therefore make the following conjecture:

**Conjecture 5.18.** *Every vertex of $Q_\ell/H^1$ takes values in $\{-1/2, 0, 1/2\}$ on every tripod of length $\ell - 1$.*

This does not follow from the proof of the $\ell = 2$ case because for higher lengths, the tripods interact in more complicated ways; however, the proof should be similar.

*Example* 5.19 ($Q_3$ in rank 2). For rank 2, the unit ball in $Q_3/H^1$ is an 11 dimensional polyhedron with 168 faces, and 4356 vertices of which 6 correspond to the vertices of the unit ball in $Q_2/H^1$ (i.e. to the rotation quasimorphisms associated to basic realizations). There are 216 tripods with edges of length 2, so 48 of the tripod faces are not of maximal dimension.

5.6. **Generalizations.** One natural example to consider is the generalization from groups to groupoids. Groupoid generators for free groups arise naturally e.g. by passing to finite index subgroups, since a nontrivial covering space of a graph will necessarily have more than one vertex.

*Example* 5.20 (Groupoid generators). In the sequel, especially in § 7, we will be interested in more complicated graphs $X$ with $\pi_1(X) = F$. Such a graph determines in an obvious way groupoid generators for $F$, and it makes sense to define counting quasimorphisms with respect to such groupoid generators. We may define fatgraphs over $X$, and surface realizations compatible with a given graph structure (see § 7 for precise definitions), and the analog of Theorem 5.11 holds in this context with essentially the same proof, namely that the codimension one faces of the unit ball in $Q_2/H^1$ are exactly the tripod faces $\pi_{x,y,z}$ associated to a cyclic ordering on a triple of edges incident to some vertex of $X$. Moreover, if $|v|$ denotes the valence of a vertex $v$ of $X$, the product $\oplus_v S_{|v|}$ of symmetric groups acts by permutation on the tripod faces, and extends to a linear action on $Q_2/H^1$. Of course, for each $v$ the action of $S_{|v|}$ factors through the induced action on the set of cyclic orders on triples; so when $|v| = 3$ each $S_{|v|}$ factors through $\mathbb{Z}/2\mathbb{Z}$.

## 6. Stallings traintracks and surface subgroups of doubles

The virtue of working with traintracks is to replace some (typically infinite dimensional) subcone of $B^+$ with a finite dimensional approximation $W^+$. By judicious choice of traintracks, this lets us reduce certain *a priori* infinite optimization problems to finite linear programming.

In this section we discuss optimization problems associated to a pair $(F, \{F_i\})$ where $F$ and each $F_i$ is a finitely generated free group, and each $F_i \subset F$. The main application is the Double Norm Theorem, which says that if $G$ is obtained by doubling a free group $F$ along a collection of finitely generated subgroups, then the unit ball in the Gromov norm on $H_2(G; \mathbb{R})$ is a rational polyhedron, and every rational class is projectively represented by a sum of closed surface subgroups.

### 6.1. **Stallings traintracks.**

Let $F$ be a finitely generated free group, and $F'$ a finitely generated subgroup of $F$. If we fix a free generating set for $F$, then the generators of $F'$ can be expressed as reduced words in $F$.

Starting with a (pointed) bouquet of circles whose (oriented) edges are labeled by reduced generators for $F'$, Stallings [36] defined a *folding* procedure which produces a new (finite) pointed graph $\Gamma$, with (oriented) edges labeled by reduced words in $F$, in such a way that non-backtracking walks in $\Gamma$ starting and ending at the basepoint are in bijection with elements of $F'$, where the element is obtained from the walk by reading the edge labels (taking inverse labels when the orientation of the walk disagrees with the orientation of the edges). It is useful to subdivide edges if necessary so that each edge is labeled by a single generator of $F$, and we adhere to this convention.

It is a straightforward and well-known observation that for such a Stallings folded graph $\Gamma$, there is a surjective map from the set of non-backtracking *loops* in $\Gamma$ to the set of *conjugacy* classes in $F$ that are conjugate into $F'$, where the surjection is obtained by reading edge labels as above. This map is *not* necessarily a bijection, since there might be (homotopically) distinct loops in $\Gamma$ with the same labels in $F$. This reflects the fact that there might be distinct conjugacy classes in $F'$ which become conjugate in $F$.

There are many canonical ways to replace the graph $\Gamma$ by a (directed) traintrack.

**Definition 6.1.** Let $F'$ be a finitely generated subgroup of $F$, and let $\Gamma$ be a Stallings folded graph for $F'$. The *Stallings traintrack* $T_\Gamma$ is defined as follows.

The vertices of $T_\Gamma$ are in bijection with the non-backtracking paths in $\Gamma$ of length 2, and the (directed) edges of $T_\Gamma$ are in bijection with the non-backtracking paths in $\Gamma$ of length 3. The endpoints of an edge corresponding to a path of length 3 are the initial and terminal subpaths of length 2.

The label on an edge is the label in $\Gamma$ associated to the middle segment of a path of length 3.

**Lemma 6.2.** *There is a natural bijection between non-backtracking loops in $\Gamma$ and directed loops in $T_\Gamma$.*

*Proof.* Each non-backtracking loop in $\Gamma$ determines a loop in $T_\Gamma$ by (cyclically) concatenating the successive subpaths of length 3. Conversely, each loop in $T_\Gamma$ determines a loop in $\Gamma$ by taking each edge of $T_\Gamma$ to the middle segment of the associated path in $\Gamma$. Since the condition that a path be non-backtracking is certified by the set of subpaths of length 2, the image is non-backtracking. These maps are mutually inverse. $\square$

Now let $F_1, F_2, \cdots, F_k$ be a finite collection of finitely generated subgroups of $F$. For each $i$ we have a Stallings graph $\Gamma_i$, and we let $\Gamma = \coprod_i \Gamma_i$ denote the graph obtained by disjoint union. Form $T_\Gamma$ from $\Gamma$ as above.

We let $W^+$ denote the cone of non-negative weights on $T_\Gamma$, and $Y^+$ the cone of non-negative polygon vectors. Recall (see Definition 3.3) that $W^+ = W \cap \mathbb{R}^+[E]$ where $W = \ker \partial \oplus H : \mathbb{R}[E] \to \mathbb{R}[V] \oplus H_1(F)$. But there is also a rational linear map $H' : \ker \partial \to \oplus_i H_1(F_i)$, whose restriction to $W$ has image equal to the kernel of the inclusion map $\oplus_i H_1(F_i) \to H_1(F)$.

The map $H'$ is rational linear, and therefore we conclude the following:

**Proposition 6.3.** *For each $\alpha \in \ker(\oplus_i H_1(F_i) \to H_1(F))$ define $\mathrm{scl}_F(\alpha)$ to be the infimum of $\mathrm{scl}_F(w)$ over $w$ conjugate into $\oplus_i C^+(F_i)$ representing $\alpha$ in $\oplus_i H_1(F_i)$. The function $\mathrm{scl}_F$ is piecewise rational linear on $\ker(\oplus_i H_1(F_i) \to H_1(F))$, and for any rational $\alpha$, it is achieved on some rational $w_\alpha$, projectively bounding an extremal surface in $F$.*

*Proof.* The restriction $H' : W^+ \to \ker(\oplus_i H_1(F_i) \to H_1(F))$ is rational linear. Now apply Theorem 3.24. $\square$

### 6.2. **Doubles of free groups.**

**Definition 6.4.** Let $F$ be a a group, and $F_1, F_2, \cdots, F_k$ a finite collection of finitely generated subgroups of $F$. The *double of $F$ along the $F_i$* is the fundamental group of the space $Z$ obtained as follows. Take $K$ a $K(F,1)$, and $K_i$ a $K(F_i,1)$ for each $i$, and let $X$ be obtained by attaching the mapping cone of the map $K_i \to K$ associated to the inclusion $F_i \to F$. We let $Y$ be the disjoint union of the $K_i$ in $X$, and let $Z$ be obtained from two copies of $X$ by identifying the copies of $Y$ with each other by the identity.

We denote the double by $G := D(F, \{F_i\})$. The double is a graph of groups with associated graph consisting of two vertices joined by $k$ distinct edges. Note that $Z$ as constructed above is a $K(G,1)$, and therefore we may compute the homology of $G$ by Mayer-Vietoris. In particular, we have an exact sequence

$$0 \to H_2(Z) \to H_1(Y) \to H_1(X)$$

or in other words,

$$H_2(G) = \ker(\oplus_i H_1(F_i) \to H_1(F))$$

6.3. **The Double Norm Theorem.** From Proposition 6.3 we deduce the Double Norm Theorem:

**Theorem 6.5** (Double Norm Theorem)**.** *Let $F$ be a free group, and $F_i$ a finite collection of finitely generated subgroups. Let $G$ be the double of $F$ along the $F_i$; i.e. $G := D(F, \{F_i\})$. Then the unit ball in the Gromov norm on $H_2(G)$ is a rational polyhedron, and every rational class in $H_2(G)$ is projectively represented by a finite sum of injective closed surface subgroups.*

*Proof.* Let $A \in H_2(G)$ correspond to $\alpha \in \ker \oplus_i H_1(F_i) \to H_1(F)$. Then by Proposition 6.3, $\mathrm{scl}_F(\alpha)$ is rational, and projectively realized by some surface $S$ with boundary in $Y$. Doubling this surface gives a closed surface in $Z$ projectively representing $A$. By construction, this surface realizes the Gromov norm of $A$; in fact, $\|A\|_\infty = 4\,\mathrm{scl}_F(\alpha)$. It follows that the components are injective. The rational polyhedrality of the norm also follows from $\|A\|_\infty = 4\,\mathrm{scl}_F(\alpha)$ and Proposition 6.3. □

If we consider a more complicated amalgam like $G = F_1 *_{F_3} F_2$, then $A \in H_2(G)$ corresponds to classes $\alpha \in H_1(F_3)$ in the kernel of inclusions into both $H_1(F_1)$ and $H_1(F_2)$. We get an inequality $\|A\|_\infty \ge 2(\mathrm{scl}_{F_1}(\alpha) + \mathrm{scl}_{F_2}(\alpha))$, and we can even find extremal surfaces $S_1$ and $S_2$; however there is now no guarantee that the boundaries will glue up.

A more refined estimate gives $\|A\|_\infty \ge \inf_w 2(\mathrm{scl}_{F_1}(w) + \mathrm{scl}_{F_2}(w))$ where the infimum is now taken over the rational polyhedron of weights $w \in W^+$ representing $\alpha$ in $H_1(F_3)$ (in particular, the right hand side of this inequality is piecewise rational linear on $H_2(G)$). If the support of some infimal $w$ is an *embedded* 1-manifold in $T$, extremal surfaces on either side can be glued up (after passing to a cover), giving rise to an injective surface in $G$. We are therefore motivated to make the following conjecture:

**Conjecture 6.6** (Norm Conjecture)**.** *Let $G$ be a graph of groups whose vertex and edge groups are free. Then the unit ball in the Gromov norm on $H_2(G)$ is a rational polyhedron, and every rational class is projectively represented by a finite sum of injective closed surface subgroups.*

## 7. Endomorphisms and traintrack dynamics

Let $G$ be a group, and let $\phi : G \to G$ be an endomorphism. Pre and post composition with an inner automorphism of $G$ defines an equivalence relation on the set of endomorphisms, and we call the equivalence classes *Outer Endomorphisms.* If $x \in G$ determines the inner automorphism $I_x$, given by the formula $I_x(y) = xyx^{-1}$ for all $y \in G$, then $\phi I_x = I_{\phi(x)}\phi$, and therefore composition of Outer Endomorphisms is well-defined and associative.

Let $F$ be a free group. By abuse of notation, we denote the semigroup of Outer Endomorphisms of $F$ by $\mathrm{End}(F)$, and for the sake of brevity we frequently drop the adjective "Outer" in the sequel.

7.1. **Linearizing endomorphisms.** We are concerned in the sequel with the action of $\mathrm{End}(F)$ on the linear spaces and objects defined in the first part of this paper. There is an obvious linear action of $\mathrm{End}(F)$ on the spaces $C(F)$ and $B(F)$, but without more hypotheses there is no natural way to define $\phi : W_\ell \to W_\ell$ so that $\phi \circ \Phi_\ell(c) = \Phi_\ell \circ \phi(c)$ for all $c \in C$. However, $\phi$ is well-defined as a linear map between suitable *different* weight spaces:

**Proposition 7.1.** *Suppose $\phi$ is injective. Then for any $\ell$ there is an $\ell'$ so that there is a well-defined map $\phi : W_{\ell'} \to W_\ell$ in the sense that $\phi \circ \Phi_{\ell'}(c) = \Phi_\ell \circ \phi(c)$ for any $c \in C$.*

*Proof.* Since every finitely generated subgroup of a free group is quasiconvex, the map $\phi : F \to F$ is a quasi-isometric embedding. By the definition of quasi-isometry, and elementary properties of free groups, there is a constant $T$ so that if $u$ and $v$ are reduced, then $\phi(uv)$ is obtained from $\phi(u)$ and $\phi(v)$ by canceling at most $T$ letters from the suffix of $\phi(u)$ and the prefix of $\phi(v)$ (or else we could find points arbitrarily far apart in $F$ whose image in $\phi(F)$ is a bounded distance apart).

It follows that for any word $v$ with $\mathrm{length}(v) \geq 2T$ there is a well defined *core* $\phi_c(v)$ which is a maximal subword of $\phi(v)$ that is never canceled in the image of $\phi(uvw)$ for any reduced $uvw$. By the definition of $T$, we have an estimate $\mathrm{length}(\phi_c(v)) \geq \mathrm{length}(\phi(v)) - 2T$. Moreover if $v$ is a prefix of $v'$, then $\phi_c(v)$ is a prefix of $\phi_c(v')$.

Since $\phi$ is a quasi-isometric embedding, there is some $\ell'$ with the property that $\mathrm{length}(\phi(v)) \geq \ell + 2T - 1$ whenever $\mathrm{length}(v) \geq \ell' - 1$. Let $v$ be a word of length $\ell'$, and let $v^-$ be obtained from $v$ by removing the last letter. Notice that $\phi_c(v^-)$ is a prefix of $\phi_c(v)$ and satisfies $\mathrm{length}(\phi_c(v^-)) \geq \ell - 1$. By abuse of notation, for $v \in W_{\ell'}$ we define $\phi(v) \in W_\ell$ to be the sum of the (possibly empty) set of subwords of $\phi_c(v)$ of length $\ell$ whose last letter is in the suffix $\phi_c(v) - \phi_c(v^-)$.

Now suppose $w$ is a cyclic word, and let $w_i$ be successive subwords of length $\ell'$. The cores $\phi_c(w_i^-)$ wrap monotonically around the cyclically reduced representative of $\phi(w)$, and $\phi_c(w_i)$ has $\phi_c(w_i^-)$ as a prefix and $\phi_c(w_{i+1}^-)$ as a suffix. Since each $\phi_c(w_i^-)$ has length at least $\ell - 1$, every subword of $\phi(w)$ of length $\ell$ occurs in exactly one $\phi_c(w_i)$ in such a way that its last letter is in $\phi_c(w_i) - \phi_c(w_i^-)$. In other words, $\phi \circ \Phi_{\ell'}(w) = \Phi_\ell \circ \phi(w)$ as desired. □

It follows that $\phi$ induces $\phi^* : Q_\ell \to Q_{\ell'}$, and thereby an endomorphism of $Q_{\mathrm{count}}$ to itself.

*Remark* 7.2. Once we have seen that $\phi : W_{\ell'} \to W_\ell$ as in Proposition 7.1 is well-defined, it follows that it is constant on the *Outer Endomorphism* class of $\phi$, since $W_\ell$ is spanned by terms of the form $\Phi_{\ell'}(c)$ for $c \in C$, and conjugation acts trivially on $C$.

*Example* 7.3. The hypothesis that $\phi$ is injective is essential in Proposition 7.1. For example, let $F = \langle x, y, z \rangle$ and define an endomorphism by $\phi(x) = x$, $\phi(y) = y$, $\phi(z) = \mathrm{id}$. Define two cyclic words

$$w_{1,n} := xz^n yz^n xz^n yz^n \text{ and } w_{2,n} := xz^n xz^n yz^n yz^n$$

Then $\Phi_\ell(w_{1,n}) = \Phi_\ell(w_{2,n})$ for all $\ell < n+2$, but $\Phi_2(\phi(w_{1,n})) \neq \Phi_2(\phi(w_{2,n}))$.

Suppose there is some $\ell$ such that $\mathrm{length}(\phi_c(v)) \geq \mathrm{length}(v)$ for all $v$ with $\mathrm{length}(v) \geq \ell - 1$. Then the proof of Proposition 7.1 shows that $\phi : W_\ell \to W_\ell$

is well-defined. Automorphisms (with almost trivial exceptions) never satisfy this condition for any $\ell$. By contrast, this condition is *generic* (in a strong sense) for *endomorphisms*.

**Definition 7.4.** Fix a free group $F$ of finite rank, and fix a free generating set. A *random endomorphism of $F$ of length $n$* is the endomorphism determined by sending each of the generators to a random reduced word in $F$ of length $n$, where the words are chosen independently and with the uniform distribution.

**Lemma 7.5.** *Fix a free generating set $S$ for $F$, and for a given $n$ let $\phi : F \to F$ take each generator to a random reduced word of length $\le n$. Then there is a constant $C > 1$ so that with probability $1 - O(C^{-n})$ we have $\text{length}(\phi_c(x)) \ge 1$ for each generator $x \in S$, and therefore $\phi$ has a well-defined action on $W_\ell$ for all $\ell \ge 2$.*

*Proof.* Two random words of length $n$ have a common maximal prefix of order $O(\log n)$, with probability $1 - O(C^{-n})$. Since the cardinality of $S$ is fixed, it follows that $\text{length}(\phi_c(x)) \ge n - O(\log n) \ge 1$ with probability $1 - O(C^{-n})$, and therefore $\phi$ acts on $W_\ell$ for $\ell \ge 2$ by the argument of Proposition 7.1. □

**Theorem 7.6** (Random Isometry)**.** *Fix a free generating set for $F$ and let $\phi$ be a random endomorphism of $F$ of length $n$. There is a constant $C > 1$ so that with probability $1 - O(C^{-n})$, the map $\phi : W_\ell \to W_\ell$ is an* eventual isometry*. That is, there is an $m$ (depending on $\ell$ but not on $\phi$ or $n$) so that $\|\phi^{m+N}(w)\|_\ell = \|\phi^m(w)\|_\ell$ for all $w \in W_\ell$ and all $N \ge 0$.*

*Proof.* Suppose that $x, y, z$ are the labels on a tripod of length $\ell - 1$. Replace the edge labels with $\phi(x), \phi(y), \phi(z)$; the resulting graph can be (Stallings) folded, giving rise to a new tripod with edges of length $\ell(n - O(\log n))$. There are finitely many tripods of length $\ell - 1$, and therefore the action of $\phi$ is eventually periodic, acting by permutations on some finite subset of the set of tripods of length $\ell - 1$.

This collection of tripods defines a collection of tripod hyperplanes, which cut out a polyhedron in $Q_\ell/H^1$; this polyhedron is dual to a polyhedron in $W_\ell$ which is the unit ball in a (pseudo)-norm invariant under $\phi$, and whose value on $w$ is the eventually constant value of $\|\phi^m(w)\|_\ell$. □

*Remark* 7.7. Theorem 7.6 implies that the spectrum of a random $\phi$ on $W_\ell$ (where it is defined) consists of zeros and roots of unity. This theorem should be compared with Theorem 3.16 from [13] which says that a random endomorphism of $F$ is an isometry of scl on $B/\epsilon$.

The following example shows that the action on $W_2$ does not need to be injective.

*Example* 7.8. Let $F = \langle x, y\rangle$, and define $\phi : F \to F$ on generators by $\phi(x) = xY$ and $\phi(y) = yx$. We compute $\phi_c(x) = x$, $\phi_c(X) = X$, $\phi_c(y) = x$, and $\phi_c(Y) = X$. In this example, for any $v$, the core $\phi_c(v)$ is obtained from the reduced word $\phi(v)$ by removing an initial $y$ or terminal $Y$ if there is one, so for example,

$$\phi(yyXYx) = yxyxyXXYxY \text{ and } \phi_c(yyXYx) = xyxyXXYx$$

The action of $\phi$ on $W_2$ is given in coordinates by the following table:

| | |
|---|---|
| $xx$ | $xY + Yx$ |
| $xy$ | $xx$ |
| $xY$ | $xY + YX$ |
| $yx$ | $xx$ |
| $yX$ | $xy + yX$ |
| $yy$ | $xy + yx$ |
| $XX$ | $Xy + yX$ |
| $Xy$ | $Xy + yx$ |
| $XY$ | $XX$ |
| $Yx$ | $XY + Yx$ |
| $YX$ | $XX$ |
| $YY$ | $XY + YX$ |

There are 8 length 1 tripods in $F$. We denote a tripod by its outgoing labels, in cyclic order. The action of $\phi$ takes $X, Y, y \to X, Y, x$ and $y, Y, x \to y, X, x$. It interchanges $X, Y, x$ and $y, X, x$; there is a similar action on the four tripods obtained by reversing the cyclic orders on these triples.

7.2. **Graph maps.** The verbal traintracks relate naturally to the structure of $F$ as a group with a fixed generating set. When studying endomorphisms of $F$ geometrically, it is natural to think of $F$ as the fundamental group of a graph, possibly with more than one basepoint. Edges of such a graph become groupoid "generators" for $F$, and it is convenient to generalize the theory of verbal traintracks to this setting. This generalization is routine, and leads to no unpleasant surprises.

**Definition 7.9.** Let $F$ be a free group. A *graph realization* consists of a connected graph $X$ without 1-valent vertices, together with a conjugacy class of isomorphism $F \to \pi_1(X)$.

An outer endomorphism $\phi : F \to F$ induces a well-defined homotopy class of self-map $X \to X$, and conversely. The following terminology is standard:

**Definition 7.10.** Let $X$ be a graph without 1-valent vertices. A *graph map* is a map $f : X \to X$ taking vertices to vertices, and edges to (possibly empty) simplicial paths without backtracking. A *traintrack map* is a graph map $f : X \to X$ such that $f^n$ is a graph map for every positive $n$. A *graph immersion* is a graph map $f : X \to X$ which restricts to an embedding on the link of each vertex.

Notice that a graph immersion is a traintrack map.

If $f : X \to X$ is a graph map, and $\gamma$ is an immersed path in $X$, then we can define $f_i(\gamma)$ to be the immersed path (i.e. the geodesic) in $X$ in the relative homotopy class of $f(\gamma)$. Note that it makes sense to define the core $f_c(\gamma)$ (in the sense of Proposition 7.1) to be the maximal subpath of $f_i(\gamma)$ which is contained in $f_i(\gamma')$ whenever $\gamma$ is extended to an immersed path $\gamma'$. If $f : X \to X$ is a graph immersion, then $f_c(\gamma) = f_i(\gamma) = f(\gamma)$ for every immersed path $\gamma$.

**Definition 7.11.** If $X$ is a simplicial graph, we define $E_\ell^X$ to be the set of non-backtracking directed simplicial paths in $X$ of length $\ell$. We define $V_\ell^X$ to be the set of non-backtracking paths of length $\ell - 1$. In a manner completely analogous to Definition 4.1, we define $\partial : \mathbb{R}[E_\ell^X] \to \mathbb{R}[V_\ell^X]$ and $h : \mathbb{R}[E_\ell^X] \to H_1(F)$, although here the function $h$ is slightly more complicated. In order to assign a homology class to a path, we pick a spanning tree in $X$; the remaining edges give a set of generators of $\pi_1(X)$, which are each in turn associated with a homology class in $H_1(F)$. To a path that starts on one of these edges, we assign the associated homology class. The other paths are sent to zero.

Finally, we set $W_\ell^X = \ker[\partial \oplus h]$. This is the *weight space* on the set of *verbal paths* in $X$.

*Example* 7.12. Let $X$ be the standard rose (i.e. a wedge of circles), with a generator of $F$ associated with each circle in the identification $F = \pi_1(X)$. Then $W_\ell^X$ is $W_\ell$, the previously defined weight space on words of length $\ell$.

Recall the cone $B^+$. The maps $\Phi_\ell : B^+ \to W_\ell$ are defined by counting cyclic subwords. For a graph realization $X$ of $F$, we get similar maps $\Phi_\ell^X : B^+ \to W_\ell^X$ where $\Phi_\ell^X(C)$ is the collection of subpaths of the non-backtracking representative of $C$ in $X$.

*Example* 7.13. For $X$ the standard rose, $\Phi_\ell^X = \Phi_\ell$.

**Lemma 7.14.** *For $b \in B^+$ and for $X$ a graph realization of $F$ there is an inequality $\mathrm{scl}(\Phi_\ell^X(b)) \le \mathrm{scl}(b)$. Furthermore, if every primitive word in the support of $b$ has length at most $\ell - 1$, then $\mathrm{scl}(\Phi_\ell^X(b)) = \mathrm{scl}(b)$.*

*Proof.* This is completely analogous to the discussion in Section 4.3. □

### 7.3. Irreducible endomorphisms.

**Definition 7.15.** An endomorphism $\phi$ of $F$ is *irreducible* if and only if there is no proper free factor $F'$ of $F$ for which $\phi^n(F')$ is conjugate into $F'$ for some positive $n$.

Note that irreducible endomorphisms are injective.

Reducibility (i.e. failure to be irreducible) can be detected at the level of a graph map $f : X \to X$ representing $\phi$, namely that there should exist such a graph map with no invariant forest whose transition matrix $N$ fails to be irreducible, in the sense that there are $i, j$ such that $(N^n)_{i,j} = 0$ for all $n$ (see [4], Lem. 1.2).

Irreducibility is a very natural condition to impose on an endomorphism. Irreducible automorphisms can be very complicated; on the other hand, Reynolds showed that an endomorphism which is irreducible but not surjective is represented by a graph immersion:

**Theorem 7.16** (Reynolds [31], Corollary 5.5)**.** *Let $\phi : F \to F$ be an irreducible endomorphism which is not an automorphism. Then $\phi$ is represented by a graph immersion $f : X \to X$.*

**Corollary 7.17.** *Let $\phi : F \to F$ be an irreducible endomorphism which is not an automorphism. Then there is a graph realization of $\phi$ as $f : X \to X$ so that $\phi$ acts naturally on $W_\ell^X$ for all $\ell$.*

*Proof.* This is just the observation that a graph immersion (whose existence is guaranteed by Theorem 7.16) satisfies $f_c(\gamma) = f(\gamma)$ for all immersed paths $\gamma$, and therefore has $\mathrm{length}(f_c(\gamma)) \ge \mathrm{length}(\gamma)$ for all $\gamma$. Now apply the argument of Proposition 7.1. □

### 7.4. Surface realizations.

**Definition 7.18.** Let $F$ be a free group. A *surface realization* consists of a compact connected oriented surface $\Sigma$, together with a conjugacy class of isomorphism $F \to \pi_1(\Sigma)$.

By abuse of notation, we will often let $\Sigma$ stand for the realization, although the data of the conjugacy class of isomorphism (i.e. the "marking" in the language of Teichmüller theory) is an essential part of the data.

*Example* 7.19. The basic realizations obtained from a fatgraph structure on a rose $X$ defined in § 5.3 are examples of surface realizations.

A choice of basis for $F$ determines a graph realization whose underlying graph is a rose. Just as basic realizations are obtained from a choice of fatgraph structure on a rose, surface realizations may be obtained from graph realizations by a choice of fattening. Recall the definition (Definition 3.17) of a fatgraph.

A fatgraph structure on a graph $X$ determines a canonical "fattening" of $X$ to a surface $\Sigma(X)$ which contains $X$ as a deformation retract.

**Definition 7.20.** If $X$ is a graph realization of $F$, then a fatgraph structure on $X$ determines a surface realization $\Sigma(X)$ of $F$. We say the surface realization is obtained from the graph realization by *fattening*.

Let $\Sigma$ be a surface realization of $F$. An outer endomorphism $\phi : F \to F$ determines a homotopy class of self-map $\Sigma \to \Sigma$, and conversely. If $X$ is a rose, edges of $X$ correspond to a generating set (i.e. a "basis") for $F$, and fattenings of $X$ correspond to a choice of cyclic order on the symmetrized generating set.

**Definition 7.21.** An endomorphism $\phi : F \to F$ is *geometric* with respect to a surface realization $\Sigma$ if there is a hyperbolic structure on $\Sigma$ with geodesic boundary such that the induced homotopy class $\Sigma \to \Sigma$ is represented by an orientation-preserving immersion taking $\partial\Sigma$ to a collection of geodesics.

It is convenient to use the language of pleated surfaces; see e.g. [38], § 8.8–10 for an introduction. If we fix a hyperbolic structure on $\Sigma$ and a (suitable) ideal triangulation of $\Sigma$, there is a pleated representative $\Sigma \to \Sigma$ which takes each ideal triangle $\Delta$ in the triangulation to an ideal triangle. An endomorphism is geometric (in the sense of Definition 7.21) if any (equivalently all) of these pleated representatives is an immersion — i.e. if it is orientation-preserving on each ideal triangle.

*Example* 7.22. Let $\Sigma$ be a surface realization of $F$. An *automorphism* $\phi \in \mathrm{Out}(F)$ is geometric with respect to $\Sigma$ if and only if it is an element of the mapping class group $\mathrm{Mod}(\Sigma)$.

Fix a finite graph $X$. Let $\mathrm{Fat}(X)$ denote the set of fattenings of $X$. This is a finite set. Fattenings pull back under immersions; hence a graph immersion $f : X \to X$ induces an endomorphism $f^*$ of $\mathrm{Fat}(X)$.

**Lemma 7.23.** *Let $\Sigma(X)$ be a fattening of $X$ invariant under $f : X \to X$. Then the induced homotopy class of map $f : \Sigma(X) \to \Sigma(X)$ is represented by a pleated immersion.*

*Proof.* We abbreviate $\Sigma(X)$ by $\Sigma$ in the sequel.

A fattening determines a planar embedding of the universal cover $\widetilde{X}$ of $X$, and thereby a circular order on the set of ends of $\widetilde{X}$. Pick a hyperbolic structure on $\Sigma$ and an ideal triangulation. Each ideal triangle determines three ends of $\widetilde{X}$, and its orientation is determined by the cyclic order on this triple of ends. Since the fattening is invariant by hypothesis, a pleated representative takes each ideal triangle to an ideal triangle with the same orientation. Such a map is necessarily an immersion. ☐

**Theorem 7.24.** *Let $\phi : F \to F$ be an irreducible endomorphism which is not an automorphism. There is a finite power $\phi^n$ which is geometric with respect to some surface realization of $F$.*

*Proof.* By Theorem 7.16 there is a graph realization $X$ so that the induced map $f : X \to X$ is a graph immersion. Since $\text{Fat}(X)$ is finite, the map $f^*$ is pre-periodic, and therefore some power $f^n$ fixes a fattening. This fattening determines a surface realization with respect to which $\phi^n$ is geometric, by Lemma 7.23. ☐

7.5. **Skinning maps.** A pleated immersion $f : \Sigma \to \Sigma$ pulls back a hyperbolic structure, and thereby induces a *skinning map* $f^* : \text{Teich}(\Sigma) \to \text{Teich}(\Sigma)$. If $f$ is irreducible this map has no fixed point, since (for any fixed hyperbolic structure on $\Sigma$) the length of the geodesic representative of $f^n(\gamma)$ is proper for each nontrivial homotopy class of loop $\gamma$.

Nevertheless, $f^*$ will fix some projective length function $\ell$ on conjugacy classes in $F$, corresponding to some degenerating sequence of hyperbolic structures on $\Sigma$. Since the length of the boundary geodesics increases without bound, most of the boundary is contained in the thin part. The thick-thin decomposition gives $\Sigma$ the structure of a graph $X$ in an $f$-invariant way, where edges correspond to long thin rectangles with two opposite edges on $\partial\Sigma$. By irreducibility, each of the vertices of the graph corresponds to a simply-connected component of $\Sigma$. Thus the graph is homotopy equivalent to $\Sigma$, and we recover the original graph immersion $f : X \to X$ in the limit.

7.6. **Finite covers.** Let $f : X \to X$ be a graph immersion. Then there is a power $f^n$ which fixes the stars of finitely many vertices pointwise, and takes the star of every other vertex into one of the fixed stars. An invariant fattening for $f^n$ is the same thing as a choice of cyclic ordering on the edges at each fixed star.

If $X' \to X$ is a finite cover, then $f$ lifts to $f' : X' \to X'$ if and only if $f_*(\pi_1(X'))$ is conjugate into $\pi_1(X')$. This is guaranteed, for example, if $\pi_1(X')$ is a characteristic subgroup of $\pi_1(X)$. If $G$ is a finitely generated group, a natural class of finite index characteristic subgroups is given by fixing an $n$ and defining $G_n$ to be the intersection of the kernels of all homomorphisms from $G$ to $S_n$, the symmetric group of order $n$. If $G$ is residually finite, $\bigcap_n G_n = \text{id}$. The lift of a graph immersion will be a graph immersion. In this way we can construct many fattenings of finite covers of $X$ invariant under suitable powers of lifts of $f$.

*Remark* 7.25. Even if $f : X \to X$ is irreducible, it is *not* typically true that a lift $f' : X' \to X'$ will be irreducible. For example, LERF for free groups implies that $f_*(\pi_1(X))$ is a free factor of $\pi_1(X')$ for some finite cover $X'$ of $X$; the induced map on $X'$ is evidently reducible, taking $f_*(\pi_1(X))$ into itself.

## 8. Folded surfaces and $f$-folded surfaces

Given a free group and an injective self-endomorphism, we can form the associated HNN extension. Such extensions are "generically" hyperbolic and 1-ended, and we can ask which ones contain surface subgroups. In this section we introduce a class of surfaces in such extensions — the so-called $f$-folded surfaces — satisfying certain combinatorial conditions that ensure they are injective.

The first main result in this section is the Folded Feasibility Theorem 8.7, proved in § 8.2, which says that there is a fast, practical algorithm to search for $f$-folded surfaces. The algorithm is essentially a linear programming problem in a polygon space, very much like the algorithm implicit in the proof of the Traintrack Rationality Theorem 3.24, except that we are only interested in the *feasibility* of the problem, rather than finding a solution minimizing some linear functional.

The second main result is the Random $f$-folded Surface Theorem, proved in § 8.3, which shows that for a *random* endomorphism of a free group there exists an $f$-folded surface with probability 1 (in a suitable sense).

The third main result is the construction of an explicit example of an injective surface in Sapir's group, i.e. the extension of $F_2 := \langle a, b\rangle$ associated to the endomorphism $\phi : a \to ab, b \to ba$. This example was found by implementing the algorithm described in the Feasibility Theorem, and illustrates its practical utility.

We also include the results of some substantial computer experiments, which (together with the Random $f$-folded Surface Theorem) constitute some evidence to support the conjecture that every one-ended hyperbolic expanding HNN extension of a free group contains many surface subgroups.

### 8.1. HNN extensions.

**Definition 8.1.** Let $G$ be a group and $\phi : G \to G$ a homomorphism. The HNN extension (denoted $G*_\phi$) is the group given by the presentation

$$\langle G, t \,|\, tgt^{-1} = \phi(g) \text{ for all } g \in G\rangle\rangle$$

There is a canonical (from its structure as an HNN extension) surjection of $G_\phi$ onto $\mathbb{Z}$, sending $G$ to 0 and $t$ to 1. The kernel is normally generated by $G$.

Geometrically, $\phi$ induces a homotopy class of self-map from a $K(G, 1)$ to itself (by abuse of notation we denote this map also by $\phi$), and $K(G_\phi, 1)$ may be obtained as the mapping torus:

$$K(G_\phi, 1) := K(G, 1) \times [0, 1]/(x, 1) \sim (\phi(x), 0)$$

In the remainder of this section we will be interested in the case that $\phi$ is an injective homomorphism from a free group $F$ to itself which is irreducible but not surjective.

### 8.2. $f$-folded surfaces.

It turns out that the linear programming techniques from § 3 can be adapted to search for surfaces which are *injective* without necessarily being extremal. Since we are focused on the computer implementation of this method, we spell out the combinatorial details.

Let's assume that we have chosen a graph realization $X$ for which $\phi$ is represented by a graph immersion $f : X \to X$ (recall that the existence of $X$ is guaranteed if $\phi$ is irreducible but not surjective by Theorem 7.16).

**Definition 8.2.** A *fatgraph $Y$ over $X$* is a graph together with a map $\pi : Y \to X$ sending edges to edges, together with a fattening $Y \to \Sigma(Y)$ so that the induced simplicial map $\pi : \partial\Sigma(Y) \to X$ is totally geodesic (i.e. it is a simplicial immersion of a 1-manifold into $X$).

A fatgraph $Y$ is *folded* if the map to $X$ is an immersion.

If $X$ is the standard rose for $F$, a fatgraph over $X$ is the usual kind of fatgraph studied in [8], and $Y$ is folded if it is Stallings folded in the usual sense. Stallings' argument implies in general that if $Y$ is folded (for an arbitrary graph realization $X$) the induced map $\pi_1(Y) \to \pi_1(X) \cong F$ is injective. Observe that the property of being folded ignores the fat structure on $Y$, and is inherited by finite covers. Informally, we think of a fatgraph $Y$ over $X$ as a fatgraph whose edges have been labeled by edges of $X$.

If $\pi : Y \to X$ is a fatgraph over $X$, then $f(\pi) : f(Y) \to X$ is the fatgraph over $X$ with the same underlying topological space as $Y$, so that the map $f(\pi)$ is defined by the tautological formula $f(\pi) = f \circ \pi$, and $f(Y)$ is subdivided so that this map takes edges to edges. Informally, $f(Y)$ is obtained from $Y$ by replacing edge labels by their images under $f$.

**Definition 8.3.** An *$f$-fatgraph $Y$ over $X$* is a fatgraph $Y$ over $X$ together with a decomposition of the boundary of the fattening $\partial\Sigma(Y)$ into submanifolds $\partial^-$ and $\partial^+$ (each a union of components) so that there is an orientation-reversing homeomorphism $\hat{f} : \partial^- \to \partial^+$ lifting $f$ (i.e. for which $\pi\hat{f} = f\pi$).

Given a geodesic 1-manifold $\delta$ immersed in $X$, we can ask for an $f$-fatgraph $Y$ such that there is an oriented covering $\partial^- \to \delta$ compatible with the immersions to $X$. Informally we say that $Y$ "virtually bounds" $\delta + f(\delta^{-1})$.

An $f$-fatgraph $Y$ (over $X$) is a fatgraph whose boundary components are paired by $f$. We can use this pairing to glue $Y$ to $f(Y)$ by identifying the image of $\partial^+(\Sigma(Y))$ to the image of $\partial^-(\Sigma(f(Y)))$. Informally, we denote the result by $Y_1 := Y \cup f(Y)$, and similarly define $Y_n := Y \cup f(Y) \cup \cdots \cup f^n(Y)$.

Similarly we can use $\hat{f}$ to glue $\Sigma(Y)$ to $\Sigma(f(Y))$ along certain boundary components, and denote the resulting surface by $\Sigma_1(Y) := \Sigma(Y) \cup \Sigma(f(Y))$, and similarly define $\Sigma_n(Y) := \Sigma(Y) \cup \Sigma(f(Y)) \cup \cdots \cup \Sigma(f^n(Y))$.

Notice that there are two surfaces associated to $Y_n$, namely $\Sigma_n(Y)$ (defined as above), and the thickening $\Sigma(Y_n)$. The next lemma gives a simple condition which guarantees that these surfaces are "the same":

**Lemma 8.4.** *Suppose that no vertex of $Y$ is associated to more than one vertex in $\partial^-$. Then $\Sigma_n(Y)$ is homeomorphic to $\Sigma(Y_n)$.*

*Proof.* Each $\Sigma(f^i(Y))$ deformation retracts to $f^i(Y)$, and the tracks of this deformation are proper essential arcs which deformation retract to points in the edges of $Y$, and proper essential trees which deformation retract to the vertices of $Y$. Glue up the tracks of the deformation of each $\Sigma(f^i(Y))$ to the tracks in each $\Sigma(f^{i+1}(Y))$ by the identification on the boundary. The result is a decomposition of $\Sigma_n(Y)$ into graphs, such that the space obtained by quotienting each graph to a point is $Y_n$. We need to check that each such graph is a tree, so that the quotient map is a homotopy equivalence, and therefore that $\Sigma_n(Y)$ is homeomorphic to $\Sigma(Y_n)$. But the hypothesis that no vertex of $Y$ is associated to more than one vertex in $\partial^-$ implies that every track $\tau$ in each $\Sigma(f^i(Y))$ is attached by at most one point (i.e. a point of $\hat{f}^i(\partial^-)$) to $\Sigma(f^{i-1}(Y))$, and consequently each such track $\tau$ deformation retracts in a canonical way to this point. By induction, each component of the decomposition of $\Sigma_n(Y)$ deformation retracts to a single track in some $\Sigma(f^i(Y))$ (the least $i$ for which the intersection of the component with $\Sigma(f^i(Y))$ is nonempty) and the conclusion follows. □

Now for any $f$-fatgraph $Y$ we distinguish, amongst the vertices of $\partial^+$, those which are in the image of vertices of $\partial^-$ under $f$, and call these *$f$-vertices*.

**Definition 8.5.** An $f$-fatgraph $Y$ over $X$ is *$f$-folded* if it satisfies the following conditions:

1. as a graph over $X$ (forgetting the fat structure), the graph $Y$ is Stallings folded;
2. every $f$-vertex in $\partial^+$ is associated to a 2-valent vertex of $Y$;
3. no vertex of $Y$ is associated to more than one $f$-vertex in $\partial^+$; and
4. no vertex of $Y$ is associated to more than one vertex in $\partial^-$.

**Theorem 8.6.** *Suppose $Y$ is $f$-folded. Then the suspension of $\Sigma(Y)$ is injective in $F*_\phi$.*

*Proof.* By Lemma 8.4 and condition (4), it suffices to show that $Y_n \to X$ is folded for all $n$. But this follows by induction, since the hypotheses imply that each vertex of $Y_n$ of valence $> 2$ whose restriction to $Y_{n-1}$ has valence 2 is locally isomorphic to some vertex in $Y$. Since foldedness is local, the conclusion follows. ☐

The main theorem in this subsection is the existence of a fast, practical algorithm to find $f$-folded surfaces. The proof is straightforward, being a variation on the algorithms described in § 3. The main point is just that the property of being $f$-folded is *local*, and all $f$-folded surfaces with prescribed boundary may be built from finitely many pieces.

**Theorem 8.7** (Folded Feasibility)**.** *Given $\phi : F \to F$ and a graph realization $X$ such that $\phi$ is represented by $f : X \to X$, and given $\delta$ an immersed 1-manifold in $X$, there is a polynomial time algorithm to check whether there is an $f$-folded fatgraph $Y$ over $X$ which virtually bounds $\delta + f(\delta^{-1})$.*

*Proof.* First we show that there is a polynomial time to check whether there is a folded fatgraph $Y$ over $X$ which virtually bounds $\delta + f(\delta^{-1})$.

We define finite sets of edges $E$, arcs $A$ and polygons $P$, and linear maps $D_A : \mathbb{R}[P] \to \Lambda[A]$ and $D_E : \mathbb{R}[P] \to \mathbb{R}[E]$ as in § 3.2. By definition there is a simplicial immersion $h : \delta \cup f(\delta^{-1}) \to X$ taking oriented edges to oriented edges, which plays the role of the evaluation function. There is an involution $\iota$ on oriented edges of $X$ which acts by reversing the orientation. An *edge* is an oriented edge of $\delta + f(\delta^{-1})$. If $e, e'$ are oriented edges of $\delta + f(\delta^{-1})$ which map by $h$ to the same edge of $X$ but with opposite orientation, we say that the pairs of vertices of $e, e'$ mapping to the same vertex of $X$ are an *arc*, and all (unoriented) arcs arise in this way. Thus a rectangle $R$ (in the sense of Definition 3.12) is associated to each such pair $e, e'$, and $D_A(R) = a + a'$ (where $a, a'$ are the associated oriented arcs), and $D_E(R) = e + e'$.

A *polygon* is either a rectangle as above, or a cyclic list of arcs, satisfying the following properties. Firstly, we require that each polygon should have at least three arcs. Each arc is (for some orientation) the boundary of a rectangle. Secondly, we require that for each arc $a$ that the arc obtained by reversing the orientation is in the boundary of some rectangle. The rectangle may not be unique, but it will map by $h$ to a unique edge of $X$. Finally we require that the the edges associated to a polygon by this method are *distinct* from one another.

Let $D := D_A \oplus D_E : \mathbb{R}[P] \to \Lambda[A] \oplus \mathbb{R}[E]$, and let $y$ be a non-negative integral vector in the kernel of $D$. Then we can build a surface from rectangles and polygons by gluing them along arcs. The result is evidently a fatgraph $Y$ over $X$, and the conditions on polygons imply that the map to $X$ is an immersion. Hence the underlying graph is folded. Conversely, each such folded fatgraph over $X$ with boundary in $\delta + f(\delta^{-1})$ gives rise to a non-negative integral vector in the kernel of $D$.

To ensure that the resulting fatgraph is $f$-folded, we first to restrict the class of rectangles, so that each rectangle has at most one edge on $\delta$. Secondly, we need to restrict the class of (non-rectangle) polygons, so that of the rectangles glued to a given polygon, at most two of them have an edge on $\delta$, and these edges are adjacent. Thirdly, we need to insist that no (non-rectangle) polygon can have an arc which ends on an $f$-vertex, and no arc can both start and end at an $f$-vertex.

These conditions just restrict to explicit subsets $A' \subset A$ and $P' \subset P$, and any vector $y$ as above gives rise to an $f$-folded fatgraph. Conversely, any $f$-folded fatgraph determines a vector with polygons in $P'$ and arcs in $A'$, so we are done. □

8.3. **Random HNN extensions contain $f$-folded surfaces.** Recall (Definition 7.4) that a random endomorphism of $F$ of length $n$ is obtained by sending the generators (in a fixed generating set) to random reduced words of length $n$. We refine this definition slightly as follows:

**Definition 8.8.** A *random reduced endomorphism of* $F$ is a random endomorphism $\phi$ conditioned on the fact that $\phi$ is represented by a graph immersion of the standard rose.

We remark that there is a uniform $\epsilon > 0$ such that a random endomorphism of $F$ of length $n$ is reduced with probability $\ge \epsilon$, independent of (sufficiently large) $n$.

The main theorem of this subsection is that for any fixed $k \ge 2$, the HNN extension associated to a random endomorphism of $F_k$ of length $n$ contains an essential surface subgroup, with probability going to 1 as $n$ goes to infinity.

**Theorem 8.9** (Random $f$-folded Surface)**.** *Let $k \ge 2$ be fixed, and let $F$ be a free group of rank $k$. Let $\phi$ be a random endomorphism of $F$ of length $n$. Then the probability that $F*_\phi$ contains an essential surface subgroup is at least $1 - O(C^{-n^c})$ for some $C > 1$ and $c > 0$.*

The remainder of this subsection is devoted to the proof of Theorem 8.9.

If $\phi$ is a random *reduced* endomorphism, the essential surface subgroup can actually be taken to be $f$-folded in the sense of Definition 8.5. In the general case we must show how to generalize the definition of $f$-folded when $f$ is not necessarily an immersion, and show that the surface we construct can be taken to be $f$-folded in this new sense. For the sake of clarity we postpone the discussion of this generalization, and first prove the theorem under the hypothesis that $\phi$ is *reduced.*

Notice that we do not specify the chain $\delta$ for which $\delta + f(\delta^{-1})$ virtually bounds an $f$-folded surface. The fact is that we can taken *any* (nontrivial) homologically trivial chain for $\delta$. To see why, let $w$ be a homologically trivial cyclically reduced word, and let $v = \phi(w^{-1})$. With overwhelming probability, we can find a copy of $w^{-1}$ in $\phi(a)$ for any generator $a$ appearing in $w$, and use this to cancel $w$ in some partial fatgraph. Similarly, if $u$ is a subword of $v$ of length 2 containing an $f$-vertex $f$-vertex, we can (again with overwhelming probability) find a copy of $u^{-1}$ somewhere in $\phi(a)$, and glue these pairs of segments in our partial fatgraph. Since $w$ is fixed but $n \to \infty$, this shows that the $f$-folded condition more or less reduces just to the problem of finding a *folded* fatgraph with boundary a random homologically trivial reduced word of length $\sim n$. For the sake of simplicity however we will give the proof for the word $w = abAB$ (note our convention of writing $a^{-1}$ as $A$ and so on).

8.3.1. *Building $Y$ by inductively pairing segments.* A fatgraph $Y$ bounding $w+v$ will be obtained by decomposing $w+v$ into disjoint subwords which will be glued (along opposite sides of an edge of $Y$) in mutually inverse pairs. We will describe how to build $Y$ by inductively specifying pairs of subwords to glue. After some but not all of these subwords have been glued, we have a *partial fatgraph*, consisting of the union of glued segments, together with a collection of unglued cyclic words, which may be thought of as a chain in $B_1(F)$. In order to control the local structure of the resulting fatgraph (to ensure $f$-foldedness), we need to keep track of some extra data in the unglued cyclic words, namely the labels on edges of the (partially glued) fatgraph that have a vertex on an unglued edge. We will ensure as we glue that there is at most a single terminal edge of the partially glued fatgraph in incident to an unglued segment, and we call this terminal edge a *tag*. Colloquially, we refer to the unglued collection of words together with this data as a *chain with tags*. Thus, the problem of finding a $f$-folded fatgraph bounding a given chain can be reduced, by gluing up a pair of inverse subwords, to the problem of finding a $f$-folded fatgraph bounding a (simpler) chain with tags. Pairs of inverse segments in a chain of tags may themselves be identified, but in order to ensure the $f$-foldedness condition, we must be careful that the resulting vertices in the fatgraph satisfy the necessary conditions spelled out in Definition 8.5. The data of the labels on the tags is enough to determine this.

We introduce the following notation. If $v$ is a segment appearing somewhere in a cyclic word, we write this as $\cdot v \cdot$. We indicate a tag in the middle of a segment by $\cdot v_1 \cdot_z v_2 \cdot$, where $z$ is the letter on the tag oriented to point *into* the segment. Suppose we have segments $\cdot w_1 p v_1 \cdot$ and $\cdot v_2 P w_2 \cdot$ for inverse subwords $p, P$. Let $z_1$ be the last letter of $P$, and $z_2$ the last letter of $p$. Then in this notation, after gluing $p$ to $P$ we create segments $\cdot w_1 \cdot_{z_1} w_2 \cdot$ and $\cdot v_2 \cdot_{z_2} v_1 \cdot$.

8.3.2. *First step: reducing $f$-foldedness to foldedness.* Here and in what follows, we let $x = \phi(a)$ and $y = \phi(b)$, and use $X = x^{-1}$ and $Y = y^{-1}$. Since by hypothesis $\phi$ is represented by a graph immersion, the word $\phi(abAB)^{-1} = yxYX$ is cyclically reduced. We write $x = x_L x' X_R$ and $y = y_L y' y_R$ where $|x_L| = |x_R| = |y_L| = |y_R| = \log(n)^2$, which makes sense provided $n \gg \log(n)^2$.

There are 4 $f$-vertices in $yxYX$, one between the last letter of $y_R$ and the first letter of $x_L$, and so on. Let $\sigma$ be the segment of $yxYX$ of length 2 containing the last letter of $y_R$ and the first letter of $x_L$. With probability $1 - O(C^{-n})$ we can find a copy of $\sigma^{-1}$ in $x_L$, and we can pair these segments. We do something similar with each of the 2-letter segments containing an $f$-vertex. Finally, we look for a copy of $\cdot pbaBAP \cdot$ in $x_L$ (say), disjoint from the eight subwords paired so far, for some letter $p$, and glue the $p$ and $P$ to create a loop labeled $baBA._p$, and then glue this to the loop $abAB$. The result of this gluing is to produce a new chain with a $._P$ tag, which differs from $yxYX$ only in the segments $x_L, x_R, y_L, y_R$. It is evident that finding an $f$-folded surface bounding $abAB + yxYX$ reduces to finding a folded surface bounding this new chain with tags.

8.3.3. *Second step: folding off short loops.* We now describe a procedure to modify the subwords $y', x', Y', X'$ contained in $y, x, Y, X$ as above (note that these entire subwords survive intact to the chain with tags at this stage). Except for the constraint that $yxYX$ is cyclically reduced, it is almost true that $y'$ and $x'$ are *independent* random words of length $n - 2\log(n)^2$ in the uniform distribution. We use the following trick to ignore the cyclically reduced constraint. If $y, x$ *were* independent random reduced words of length $n$ in the uniform distribution, there would still be a uniformly positive probability ($> 5\%$ chance, say) that $yxYX$ is cyclically reduced. If we can prove that some property of $y$ and $x$ holds with probability $1 - O(C^{-n^c})$, then even after conditioning on the cyclically reduced constraint, the property will still hold with probability $1 - O(C^{-n^c})$ (at the cost of adjusting the implicit constants involved). So we are justified in arguing as though $y'$ and $x'$ are independent random words in the uniform distribution, and are therefore produced by a *stationary Markov process.*

We perform the following process. As we read off the letters of $y'$ one by one, we look for a segment of length 11 of the form $\cdot v_1 puPv_2 \cdot$, where $v_1 \neq v_2^{-1}$ and $p, P$ are letters, and $|u| = 7$. Then the $p$ and $P$ can be glued, producing the segment $\cdot v_1 \cdot_P v_2 \cdot$ and the loop $u \cdot_p$. We call $u \cdot_p$ a *short loop*, and we call this the operation of *folding off a short loop.* A short loop is determined by a cyclically reduced word $u$, a choice of a "basepoint" for $u$, and a letter $p$ not equal to the last letter of $u$ or to the inverse of the first letter of $u$. This data $(u, p)$ is called the *type* of a short loop. Let $L_{7,k}$ denote the number of types of short loop of length 7 in a free group of rank $k$, so for example $L_{7,2} = 4376$.

After folding off a short loop, we skip ahead a fixed number of letters in $y'$ (say 2 letters), and then examine the segments of length 11 one by one until we get another chance to fold off a short loop. If we fold off every short loop that we can by this process, we see that different types of short loops will be produced with different probabilities, but since there are only a finite number of different types of short loops, the ratio of these probabilities are bounded. For a given short loop type $\ell$, let $\mu_\ell$ be the relative probability that a short loop generated by this process is of type $\ell$, and let $\mu_0$ be the minimum of this collection of relative probabilities (note that the vector of $\mu_*$ is only well-defined up to scale). We adjust the process of folding off short loops as follows: whenever we have the opportunity to fold off a short loop of type $\ell$, we do so with probability $\mu_0/\mu_\ell$.

Explicitly, we start with $\cdot y' \cdot$ a random word generated by a stationary Markov process, and successively fold off short loops of type $\ell$ each time they appear with probability $\mu_0/\mu_\ell$, or leave them intact with probability $1 - \mu_0/\mu_\ell$, and the result of this process is a new segment $\cdot y'' \cdot$ of length $O(n)$ together with a collection $\mathcal{L}_y$ of short loops, also of cardinality $O(n)$.

Observe the following two facts. Firstly, the tagged word $y''$ is itself generated by a stationary Markov process. Secondly, by the Chernoff inequality, with probability $1 - O(C^{-n^c})$, for every type $\ell$ the number of loops of type $\ell$ is $|\mathcal{L}_y|/L_{7,k} \pm O(n^{1/2+\epsilon})$. We perform similar folding-off processes for $x', Y', X'$ to produce tagged words $\cdot x'' \cdot$, $\cdot Y'' \cdot$, $\cdot X'' \cdot$ and collections $\mathcal{L}_y, \mathcal{L}_X, \mathcal{L}_Y$. It is not important to ensure that $X'' = (x'')^{-1}$, but we can enforce this if we wish. Let's denote the union of the $\mathcal{L}_*$ by $\mathcal{L}$.

8.3.4. *Third step: random pairing of long subwords.* Notice that for any letters $v_1, v_2$ that there is some choice of $p, u$ so that $\cdot v_1 puPv_2 \cdot$ is reduced. It follows that if we fix some big constant $N$ that every reduced word of length $N$ will appear with definite frequency as a subword of $y''$ (and likewise $x'', Y'', X''$). Moreover, for any $v_1, v_2$ there are at least two possibilities for $p$ as above, and each possibility occurs with equal probability.

Let's subdivide $y''$ and $Y''$ into successive subwords whose lengths alternate between $N$ and either 100 or 101 (the latter chosen randomly and independently with equal probability). The successive (tagged) subwords of length $N$ can be thought of as being generated by an aperiodic ergodic Markov chain. For each word $\sigma$ of length $N$ we can consider the collection of possible tags that could appear, and call this a tagged copy of $\sigma$. A tagged copy of $\sigma$ and a tagged copy of $\sigma^{-1}$ are *compatible* if they can be glued in such a way that the result is (locally) folded. For the sake of definiteness, we define an involution $\iota$ on the set of tagged words so that for any tagged word $\sigma$, the words $\sigma$ and $\iota(\sigma)$ are compatible. First, the underlying word of $\iota(\sigma)$ should be $\sigma^{-1}$. Second, for each tag $\cdot v_1 \cdot_u v_2 \cdot$ we distinguish two cases. If $u = V_1$ we set $\iota(\cdot v_1 \cdot_{V_1} v_2 \cdot) = \cdot V_2 \cdot_{v_2} V_1 \cdot$. Otherwise, we set $\iota(\cdot v_1 \cdot_u v_2 \cdot) = \cdot V_2 \cdot_U V_1 \cdot$.

By construction and the discussion above, for every tagged subword $\sigma$, the expected number of copies of $\sigma$ in $y''$ is equal to the expected number of copies of $\iota(\sigma)$ in $Y''$, and similarly for $x''$ and $X''$. Again by the Chernoff inequality, we can glue compatible pairs of subwords in $y''$ and $Y''$ with $O(n^{1/2+\epsilon})$ exceptions, with probability $1 - O(C^{-n^c})$. After this gluing we are left with a tagged chain $z$ of total length approximately $100n/N$, plus the collection $\mathcal{L}$ of (almost perfectly equidistributed) short loops, which has total length $\ge C.n$ for some fixed constant $C$ (independent of $N$). If we choose $N$ so that $N \gg C$ then the total length of the short loops is much, much greater than the total length of $z$.

*Remark* 8.10. The requirement that tags must be compatible when gluing inverse words forbids us from simply gluing $y''$ to $Y''$ and $x''$ to $X''$, assuming we had arranged for $Y'' = (y'')^{-1}$ and $X'' = (x'')^{-1}$. This is why we need to be a bit careful in defining $\iota$ so that $\sigma$ and $\iota(\sigma)$ are compatible.

8.3.5. *Fourth step: building $Z$.* At the next stage we show how to partially glue up some (relatively small) subset of the short loops to produce a tagged copy of $Z$ (i.e. the chain inverse to $z$) whose tags are compatible with the tags of $z$. We can then glue $z$ to $Z$ and be left with only short loops. Since the total mass of $z$ is so much smaller than that of $\mathcal{L}$, the remaining collection of short loops will still be almost equidistributed.

The basic construction step is as follows. Suppose that we have two loops $vpqr$ and $v'RQP$ where $v$ and $v'$ are words, and $p, q, r$ are letters. We can glue $pqr$ to $RQP$ to produce the tagged loop $._r v ._P v'$. Note that for any two letters $p$ and $r$ there is some reduced word $pqr$ of length 3, so the resulting tags can be anything (subject, of course, to the word $vv'$ being cyclically reduced).

We abbreviate a word of length $d$ by the number $d$, so that a short loop may just be abbreviated by .7 (the dot preceding the 7 is the tag). A subsegment of such a word will be indicated by brackets, hence 4(2.1) indicates a short loop with a choice of specific segment of length 3 with the tag in its interior. Denote the operation of gluing by $\sqcup$ and the result by $\to$. Then, providing labels are compatible, we can perform an operation such as

$$4(2.1) \sqcup 4(2.1) \to .4.4$$

See Figure 3.

Consider the following sequence of gluings.

$$4(2.1) \sqcup 4(2.1) \to .4.4 = .3(1.2)2$$

$$.3(1.2)2 \sqcup 4(2.1) \to .3.4.2 = .1(2.1)3.2$$



FIGURE 3. The meaning of the operation $4(2.1) \sqcup 4(2.1) \to .4.4$

$$.1(2.1)3.2 \sqcup 4(2.1) \to .1.4.3.2 = .1.3(1.2)1.2$$

$$.1.3(1.2)1.2 \sqcup 4(2.1) \to .1.3.4.1.2 = .1.1(2.1)3.1.2$$

and so on. If we choose suitable short loops as the $4(2.1)$ summands at each stage, the result of $m$ gluings will be a loop of length $7 + m$ representing an arbitrary reduced word and with arbitrary tags. Since the total length of $z$ is so small compared to that of $\mathcal{L}$, we can build a copy of $Z$ from roughly $|Z|$ short loops, with tags compatible with the tags of $z$, and then glue $Z$ to $z$. We have therefore reduced to an almost equidistributed collection $\mathcal{L}'$ of short loops. We introduce the following terminology. We say that a collection $\mathcal{L}'$ of short loops (in some $F_k$) is *equidistributed with error* $\epsilon$ if for any two types $\ell, \ell'$ the ratio of the number of short loops of type $\ell$ to the number of type $\ell'$ is between $1 - \epsilon$ and $1 + \epsilon$. With this terminology, we have reduce the problem of exhibiting an $f$-folded surface to folding a collection $\mathcal{L}'$ of short loops in $F_k$ which is equidistributed with error $C/N$. By choosing $N$ sufficiently large (for fixed $C$) we may arrange for the error, which we abbreviate to $\epsilon_1$, to be as small as we like.

8.3.6. *Fifth step: linear programming.* Let $\Sigma_k$ denote the set of types of (tagged) short loops in $F_k$, and let $\mathbb{R}^+[\Sigma_k]$ denote the cone of non-negative linear combinations of elements in this set. The subspace of finite linear combinations which bound a folded surface is a closed rational convex subcone of this space, contained in the subspace of linear combinations representing 0 in homology. Call this subset the *feasible cone*. We claim that the projectivization of the feasible cone contains an open neighborhood of the uniform vector (i.e. the vector all of whose components are equal) in the projectivization of the subspace representing 0 in homology. This will show that with overwhelming probability $\mathcal{L}'$ as above is in the feasible cone, and therefore we will conclude the proof of Theorem 8.9, under the stronger hypothesis that $\phi$ is reduced.

For any fixed $k$, by Theorem 8.7 this claim reduces to finding finitely many vectors in $\mathbb{R}^+[\Sigma_k]$ whose projective hull (i.e. cone) contains the uniform vector in its interior, and testing the feasibility of a specific linear programming problem associated to each vector. In other words, it reduces to a finite computation for each $k$. But there is a trick to reduce the computation for arbitrary $k$ to the case of $k \le 8$. First, observe that for any short loop type $\ell$, a pair $\ell + \iota(\ell)$ bounds (in an obvious way), where $\iota$ is the involution defined in the third step. Now, any short loop $\ell$ in $F_k$ has at most 8 distinct letters, and therefore is contained in some copy of $F_8$. Let $v_8$ denote the uniform vector in $F_8$, and suppose that we can show feasibility for all homologically trivial chains in an open neighborhood of $v_8$ in $\mathbb{R}^+[\Sigma_8]$. Let $\tau_i : F_8 \to F_k$ denote the $k!/8!(k-8)!$ various inclusions of $F_8$ into $F_k$

determined by choosing 8 out of $k$ generators, and let $v' = \frac{8!(k-8)!}{k!}\sum_i \tau_i(v_8)$ denote the image. It is not quite true that $v'$ is equal to the uniform vector $v_k$, but both $v'$ and $v_k$ are vectors with all coordinates positive, and which are invariant under the involution $\iota$. It follows that if some open neighborhood of $v_8$ is feasible, then an open neighborhood of $v'$ is feasible, and an open neighborhood of $v' + w$ is feasible for any $\iota$-invariant non-negative vector $w$; in particular, an open neighborhood of $v_k$ is feasible. In the sequel we call this trick of adjusting a vector by an $\iota$-invariant vector the *$\iota$ trick*. Applying the $\iota$ trick to reduce from rank $k$ to rank 8 will magnify the error by a bounded amount, so we are reduced to the problem of finding a folded surface bounding a collection of short loops in $F_8$ which is equidistributed with error $\epsilon_2$, where $\epsilon_2$ depends on $\epsilon_1$ and $k$. Nevertheless, for fixed $k$, as $\epsilon_1 \to 0$ so too does $\epsilon_2 \to 0$.

However, even with this reduction, the size of the problem is far beyond what is feasible on a modern computer. We now describe how to reduce the problem from $F_8$ to $F_2$ in a series of steps, each of which magnifies the error of the equidistribution by a bounded amount.

First, we reduce from $F_8$ to $F_4$. For each loop type $\ell$ that uses at least 5 generators, the loop type $\iota(\ell)$ also uses at least 5 generators. By equidistribution, we pair as many such loops as we can, and are left with $\epsilon_2 Cn$ loops containing at least 5 generators, and $(C \pm \epsilon_2)n$ of each type of loop containing at most 4 generators, for some $C$. If $\ell$ is one of the remaining loops with at least 5 generators, write $\ell = st\cdot_u v$ where $|s| = 3$, $|t| = |v| = 2$ and $|u| = 1$. At most 4 generators appear in $t$ and $v$, so we can find a short loop of the form $V\cdot_w Txxx$ where $\cdot V\cdot_w T\cdot$ is compatible with $\cdot t\cdot_u v\cdot$, and where $x$ or $X$ is one of the 4 generators used in $t$ and $v$. By hypothesis we have many more copies of $V\cdot_w Txxx$ than of $st\cdot_u v$, and after gluing $\cdot V\cdot_w T\cdot$ to $\cdot t\cdot_u v\cdot$ we obtain a tagged loop of the form $sxxx$ which uses at most 4 generators. By the method of Step Four we may fold this $sxxx$ up with a finite collection of short loops using at most 4 generators. In this way we eliminate all loops with at least 5 vectors, and are left with loops containing at most 4 generators. This collection of loops may be partitioned equally into subsets, one for each choice of 4 out of the 8 generators. Each subset is now not equidistributed, but is equidistributed modulo an $\iota$-invariant vector, up to bounded error. So by the $\iota$ trick we may reduce to a collection of short loops which is equidistributed in $F_4$ with error $\epsilon_3$. Again, $\epsilon_3$ depends on $\epsilon_2$ in such a way that $\epsilon_3 \to 0$ as $\epsilon_2 \to 0$.

Next, we reduce from $F_4$ to $F_3$. Again, try to cancel each $\ell$ using 4 generators with a copy of $\iota(\ell)$ until we are left with only $\epsilon_3 Cn$ such loops. Let $\ell$ be a remaining loop using 4 generators, say $a, b, c, d$. Let $\ell'$ be a short loop obtained from $\ell$ by replacing each $d$ or $D$ by one of $a, A, b, B$. Then a copy of $\iota(\ell')$ will cancel most of a copy of $\ell$, leaving a collection of loops that use only the generators $a, b, d$. Cancel these loops using the method of Step Four. After another application of the $\iota$ trick we reduce to an equidistributed collection of short loops in $F_3$ with error $\epsilon_4$.

Finally, we reduce from $F_3$ to $F_2$. This step is nearly the same as the previous step. We try to cancel each $\ell$ using 3 generators with a copy of $\iota(\ell)$ until we are left with only $\epsilon_4 Cn$ such loops. Let $\ell$ be a remaining loop in the generators $a, b, c$, and let $\ell'$ be a short loop obtained from $\ell$ by replacing each consecutive string of $c$s or $C$s by a consecutive string of $a$s or $b$s etc., and then using a copy of $\iota(\ell')$ to cancel most of a copy of $\ell$. What is left over may contain all three generators $a, b, c$, but *each component* of what is left over is either of the form $c^*a^*$ or $c^*b^*$ and therefore uses at most two generators. Each component therefore may be canceled by the method of Step Four. After one final application of the $\iota$ trick we reduce to an equidistributed collection of short loops in $F_2$ with error $\epsilon_5$. This is finally within the reach of practical computation.

Determining if a point lies in the interior of the cone on a set of vectors can be phrased as a linear programming problem by using the following lemma.

**Lemma 8.11.** *Let $x, v_1, \ldots, v_k \in \mathbb{R}^n$. If the $v_i$ span $\mathbb{R}^n$ and there is an expression $x = \sum_i t_i v_i$ with $t_i > 0$ for all $i$, then $x$ lies in the interior of the cone spanned by the $v_i$.*

*Proof.* To show that $x$ lies in the interior, it suffices to show that $x$ is not contained in any supporting hyperplane. Thus, let $H$ be any supporting hyperplane for the cone spanned by the $v_i$. Because the $v_i$ span $\mathbb{R}^n$, there is some $j$ so that $v_j$ is not in $H$. We have expressed $x = \sum_i t_i v_i$ with, in particular, $t_j > 0$. Therefore, if we decompose $\mathbb{R}^n = H \oplus \text{span}\{v_j\}$, and express $x$ in this decomposition, we will find that the coefficient of $v_j$ is not zero, so $x$ is not contained in $H$. $\square$

Using Lemma 8.11, we observe that if we are given the $v_i$ as the columns of the matrix $M$, and $x$ is a column vector, then a feasible point $y$ for the problem $Ay = x$, $y \geq 1$ provides a certificate that $x$ is contained in the interior. Feasibility testing can be phrased as a linear programming problem by setting the objective function to zero. We remark that the lower bound on $y$ is arbitrary; if $x$ is in the interior, the linear program will succeed for *some* lower bound, but we don't know a priori what it is.

The fifth step of the proof therefore reduces to

1. Find a collection of vectors $V$ in the feasible cone which together span the space of homologically trivial linear combinations of $\Sigma$.
2. Show that the uniform vector lies in the cone on $V$.

Both steps require linear programming: in order to check that a vector lies in the feasible cone, we solve a linear programming problem to find a folded surface, and to show that the uniform vector lies in the cone on $V$, we solve the linear programming problem derived from Lemma 8.11.

In order to make the many linear programming problems in step 1 feasible, we need to choose *low-density* vectors; that is, vectors with a small number of nonzeros. Recall there are 4376 short loops of length 7 and rank 2, and the space of homologically trivial linear combinations has dimension 4374. We found a collection of 9626 vectors, each with 8 nonzeros, which span this 4374-dimensional space. This required solving a few tens of thousands of small linear programming problems, which was easily accomplished by `GLPK`[20], using our program `gallop`[39]. As we built this collection, we occasionally ran a much larger linear program to determine if the uniform vector was contained in the cone (not necessarily in the interior, as that is more difficult to solve in practice). Once our cone did contain the uniform vector, we ran one final linear program to verify that it lay in the interior.

Even though these latter linear programs had only a few thousand columns and a few thousand rows, they proved quite difficult in practice. Fortunately, the proprietary software package `Gurobi`[25], which offers a free academic license, was able to solve them in a few minutes. We used `Sage`[37] to facilitate many of the final steps.

8.3.7. *Sixth step: unreduced automorphisms.* We now show how to adapt our argument to the general case of a random endomorphism which is not assumed to be reduced.

**Definition 8.12.** Let $\phi : F \to F$ be an endomorphism, and let $\lambda$ be a real number satisfying $0 < \lambda < 1$. We say $\phi$ is *$\lambda$-reduced* if, for any two non-inverse letters $a, b$ (with the possibility that $a = b$) the maximal suffix of $\phi(a)$ canceling a prefix of $\phi(b)$ has length $< \lambda|\phi(a)|$.

*Remark* 8.13. Note that since we can take $B, A$ in place of $a, b$ we also deduce the length is $< \lambda|\phi(b)|$.

*Remark* 8.14. The condition of $\phi$ being 1/2-reduced implies (in the notation of § 7) that $\text{length}(\phi_c(x)) \geq 1$ for each generator $x \in S$, and therefore that $\phi$ has a well-defined action on $W_\ell$; compare with Lemma 7.5.

Now, suppose $\phi$ is 1/2-reduced. Let $\delta$ be a cyclically reduced word, let $\phi(\delta^{-1})$ denote the *unreduced* cyclic word obtained by concatenating the words obtained by applying $\phi$ to the letters of $\delta$, and let $\delta'$ denote the word obtained by reducing $\phi(\delta^{-1})$. We think of $\phi(\delta^{-1})$ as a simplicial loop, and as $\delta'$ as a smaller simplicial loop obtained by folding pairs of canceling subwords. There is a simplicial map from $\delta^{-1}$ to $\phi(\delta^{-1})$ taking each edge to a union of edges (i.e. to a copy of $\phi(a)$ for each letter $a$ and so on); the images of the vertices of $\delta^{-1}$ are what we previously called $f$-vertices.

The condition on $\phi$ implies that each vertex of $\delta'$ is in the image of *at most* one $f$-vertex, so it makes sense (by abuse of notation) to refer to these vertices of $\delta'$ as $f$-vertices. With this definition it makes sense to define an *$f$-folded fatgraph* $Y$ (over the standard rose) to be a fatgraph bounding $\delta + \delta'$ as above, and satisfying the conditions in Definition 8.5 (where $\delta = \partial^-$, $\delta' = \partial^+$, and the $f$-folded vertices are those with the new definition).

If $Y$ is an $f$-folded fatgraph, then $\phi(Y)$, the fatgraph obtained by replacing each edge $a$ of $Y$ by the corresponding word $\phi(a)$, will *not* typically be folded, but by the 1/2-reduced condition, the result of canceling maximal pairs of adjacent subwords in the image of adjacent letters will be a new fatgraph which *is* Stallings folded, in the usual sense. Call this new fatgraph $\phi'(Y)$. Then we can glue $\phi'(Y)$ to $Y$ along $\delta'$, and as before we see that the result is still Stallings folded. Thus we have proved the following analog of Theorem 8.6:

**Theorem 8.15.** *Let $\phi$ be 1/2-reduced, and suppose $Y$ is $f$-folded. Then the suspension of $\Sigma(Y)$ is injective in $F*_\phi$.*

The proof of Lemma 7.5 implies that a random endomorphism of length $n$ is $C'.\log n/n$-reduced, with probability $1 - O(C^{n^{-c}})$, for suitable constants $C, C', c$. So the method of the previous few sections applies essentially without change to produce an $f$-folded surface in this new sense. This concludes the proof of Theorem 8.9 in full generality.

## 8.4. **Sapir's group.**

**Definition 8.16.** We define *Sapir's group* $C$ to be the HNN extension of $F_2 := \langle a, b\rangle$ by the endomorphism $\phi : a \to ab, b \to ba$.

In [33], Problem. 8.1, Sapir posed explicitly the problem of determining whether $C$ contains a closed surface subgroup, and in fact conjectured (in private communication) that the answer should be negative. This group was also studied by Crisp–Sageev–Sapir and (independently) Feighn, who sought to find a surface subgroup or show that one did not exist. Because of the attention this particular question has attracted, we consider it significant that our techniques are sufficiently powerful to give a positive answer:

**Theorem 8.17.** *Sapir's group $C$ contains a closed surface subgroup of genus 31.*

*Proof.* The theorem is proved by exhibiting an explicit $f$-folded surface. Figure 4 indicates a fatgraph whose fattening has four boundary components, three of which are (conjugates of) $babaBABA$ and the fourth of which is $\phi(babaBABA^{-3})$. The blue circles mark the $babaBABA$ components. By taking a 3-fold cover we obtain a fatgraph whose fattening has six boundary components, three of which are conjugates of $babaBABA^3$, and three of which are conjugates of $\phi(babaBABA^{-3})$. In the HNN extension $F*_\phi$ we can glue these boundary components in pairs, giving a closed surface $S$ together with a map $\pi_1(S) \to F*_\phi$.

The surface is $f$-folded, and therefore the resulting map of the surface group is injective. To see this, note that the $babaBABA$ components are disjoint from each other, the underlying fatgraph is Stallings folded, and the $f$-vertices (indicated in red) are all 2-valent. ☐

*Remark* 8.18. In fact, Sapir expressed the opinion that "most" ascending HNN extensions of free groups should not contain surface subgroups, which is contradicted by the Random $f$-folded Surface Theorem 8.9. On the other hand, the probabilistic estimates involved in the proof of this theorem are only relevant for endomorphisms taking generators to very long words, and therefore Sapir's group seems to be an excellent test case.

### 8.5. **Experimental results.**

The Random $f$-folded Surface Theorem 8.9 implies that surface subgroups of HNN extensions of free groups are very common, and implies that for sufficiently large $n$ (depending on $k$) an HNN extension associated to a random endomorphism of length $n$ is virtually certain to contain one. However, the precise meaning of "sufficiently large" implicit in the argument is far beyond the reach of computer experiment. On the other hand, the construction of an $f$-folded surface subgroup of Sapir's group suggests that such subgroups are even more plentiful than our argument certifies, and may be found even for $n$ of modest size. In this section we describe experimental results for small $n$ which complement the asymptotic conclusions of Theorem 8.9.

In order to naively use the algorithm in the Folded Feasibility Theorem 8.7 to search for $f$-folded surfaces it is necessary to represent a given automorphism $\phi : F \to F$ by a graph immersion $f : X \to X$. For ease of computational efficiency, it is very convenient to consider maps for which $X$ can be taken to be the standard rose; however, we weaken the condition that $f : X \to X$ is a graph immersion to the condition that $\phi_c(x)$ is nonempty for every generator $x$, where $\phi_c$ denotes the core of $\phi$ (see Proposition 7.1 for a discussion of cores). This means precisely that whenever $xyz$ is a reduced word of length 3, there are letters from $\phi(y)$ remaining in the reduced word obtained from the product $\phi(x)\phi(y)\phi(z)$. Note that this implies that $\phi$ is well-defined on $W_\ell$ for all $\ell \geq 2$.



FIGURE 4. A fatgraph bounding $3 \cdot babaBABA + \phi^4(babaBABA^{-3})$

Now fix $F = \langle a, b\rangle$. Let $E$ denote the set of all endomorphisms of $F$ of length 3 (i.e. taking each generator to a reduced word of length 3) and satisfying the following properties:

(1) for every generator $x$ we have $|\phi_c(x)| \geq 1$; and
(2) no power of $\phi$ sends a generator to a conjugate of a power of itself.

Note that the first condition ensures that $|\phi(w)| \geq |w|$ for every conjugacy class $w$, so (by the second condition) $\phi$ can't be an automorphism. The second condition is implied by irreducibility. The cardinality of $E$ is 752. For each $\phi \in E$, we determined whether there is an $f$-folded surface bounding $C - \phi^n(C)$, for $1 \leq m \leq 4$ and for $C$ one of the chains $a + b + AB$ or $abAB$. Continuing the search to $m = 5$ is not computationally feasible. However, it is possible to perform a *randomized* search for $f$-folded surfaces in this case: for each endomorphism, we computed $C - \phi^5(C)$ and did many *random foldings*; that is, we simply glued $C - \phi^5(C)$ to itself along many compatible segments. The resulting (tagged) chain is short enough that an exhaustive search for $f$-folded surfaces can proceed. Of course this method may fail to find an $f$-folded surface bounding $C - \phi^5(C)$ even if one exists, but it is a reasonable way to continue. For the randomized search, we tried many different

| $m$ | surface found | fraction (out of 752) |
|---|---|---|
| 1 | 0 | 0 |
| 2 | 52 | 0.069 |
| 3 | 492 | 0.654 |
| 4 | 636 | 0.846 |
| $5^*$ | 676 | 0.899 |

TABLE 1. Number of $\phi \in E$ for which there is an $f$-folded surface bounding $C - \phi^m(C)$ with $C = abAB$ or $a + b + AB$. For $m = 5$ the search was randomized.

chains $C$. See Table 1 for the summary of results. Here $\phi$ has "no surface found" for some power $n$ if we could not find an $f$-folded surface bounding $C - \phi^i(C)$ for $i \leq n$ for any $C$ which we tried. So the first line of Table 1 means that we could not find any $f$-folded surfaces using the first power of any endomorphism.

In a second experiment, we vary the length $n$ of the endomorphism (i.e. the length of the words $\phi(x)$ as $x$ ranges over the generators), and compute the probability of "easily" finding an $f$-folded surface. Explicitly, for each value $5 \leq n \leq 24$ we generated approximately 5000 random endomorphisms of length $n$. For each such $\phi$, we searched for an $f$-folded surface bounding $C - \phi(C)$ for $C = [a, b]$. Note this is a very naive search; as our previous experiment shows, taking powers and a variety of chains $C$ produces far more $f$-folded surfaces. Nevertheless, we still observe that the probability of finding an $f$-folded increases quickly.



FIGURE 5. The experimental fraction of endomorphisms $\phi$ of length $n$ for which $[a, b] - \phi([a, b])$ bounds an $f$-folded surface.

On the basis of this experimental evidence, together with the theoretical results from this section and the next, we feel therefore reasonably confident in making the following conjecture:

**Conjecture 8.19.** *Let $\phi : F \to F$ be an irreducible endomorphism which is not an automorphism. Then for some power $\phi^m$ of $\phi$ and some chain $C$ there is an $f$-folded surface bounding $C - \phi^m(C)$. In particular, the HNN extension $F*_\phi$ should contain many closed surface subgroups.*

## 9. EXTREMAL SURFACES AND FLAT SURFACES

In this section we are concerned with finding surface subgroups of HNN extensions by methods more closely related to the theory of stable commutator length, and the techniques developed in § 2–4. We distinguish between four classes of injective surface subgroups in an HNN extension, related to each other by the following diagram of inclusions

$$\begin{tikzcd}
\text{flat} \arrow[r] \arrow[d] & f\text{-folded} \arrow[d] \\
\text{extremal} \arrow[r] & \text{injective}
\end{tikzcd}$$

Here we can take $f$-folded either in the narrow sense of Definition 8.5 for an immersion $f$, or in the broader sense of $f$-folded for $f$ merely $1/2$-reduced as in Definition 8.12.

The properties of $f$-foldedness and of extremality are both certified by linear programming. An $f$-folded surface carried by a fatgraph $Y$ over $X$ is *flat* if there is an $f$-invariant fattening of $X$ which pulls back under $Y \to X$ to the given fatgraph structure on $Y$.

Extremal surfaces and flat surfaces are (by definition) homologically essential; thus no HNN extension can contain such a surface unless $\phi$ fixes some vector in $H_1(F)$. We reduce the question of such surfaces, therefore to two intermediate questions. Firstly, when does an HNN extension $F*_\phi$ have virtually positive $b_2$? And secondly, when is some class in $H_2(F*_\phi)$ (assuming one exists) represented by an extremal surface?

We cannot give firm theoretical answers to these questions; however, the tools that we have developed give us the ability to test these questions *experimentally*, and to quickly *certify* the existence of extremal or flat surfaces. One interesting aspect of this story is the existence of certain homomorphisms to $\mathbb{R}$ defined on subsemigroups of $\text{End}_\alpha(F)$, the stabilizer of a homology class $\alpha \in H_1(F)$. Such homomorphisms are associated to (counting) quasimorphisms in $Q_\ell$. In the special case of rotation quasimorphisms, the associated homomorphisms are seen to be closely related to a particular summand of the *Johnson homomorphism*, defined on the Torelli subgroup of the mapping class group; see Proposition 9.7 and the related discussion in § 9.3

## 9.1. **Nontrivial homology in finite index subgroups.**

In this subsection we consider the question: given an endomorphism $\phi$ of $F$, when is there a finite index subgroup $F'$ of $F$ and a power $\phi'$ of $\phi$ taking $F'$ into $F'$ such that $\phi'$ fixes some nontrivial vector in $H_1(F')$?

If we fix $F$ free of rank $k$, the probability that a random endomorphism of length $n$ fixes a particular basis vector in $H_1$ is of order $n^{-k/2}$ for fixed $k$ and $n \to \infty$, and of order $k^{-n/2}$ for fixed $n$ and $k \to \infty$, in each case subject to the parity condition that $n$ should be odd (the precise asymptotics are obtained by Sharp; see [35]). If $F'$ is a subgroup of $F$ of index $m$, then the rank of $F'$ is $mk - m + 1$; thinking of the length of a random endomorphism as (approximately) the stretch factor, we can think of a lift of $\phi$ to $F'$ (if one exists) as an endomorphism of length of order $n$. Of course, the lift is no longer a random endomorphism, and different lifts to different $F'$ are certainly not independent. However this heuristic suggests that it should be relatively easy to find nontrivial fixed elements in $H_1$ in finite index subgroups of $F$. In fact, the probability of finding such fixed vectors should go to 1 exponentially fast in the index. At least for small values of $n$, this heuristic is borne out in practice.

Fix $F = \langle a, b\rangle$, and let $E$ denote the set of endomorphisms $\phi$ of length 3 for which no power of $\phi$ sends a generator to a conjugate of a power of itself. Then $|E| = 848$. For each $\phi \in E$, we searched for finite index subgroups $F'$ of $F$ with

$\phi(F') \subset F'$ and such that $\phi$ fixes some nontrivial vector in $H_1(F')$. We find that *in every case* there is such a finite index subgroup with index at most 10. For length 4 endomorphisms, of which 11,132 satisfy our condition, the number without such a finite index subgroup through index 11 is 328, or 2.9%, at which point continuing the computation becomes infeasible.



FIGURE 6. The fraction of length 3, left, and length 4, right, irreducible endomorphisms $\phi$ for which there is a subgroup of $F_\phi$ with finite index at most $i$ and with nontrivial $H_2$. The plot on the left reaches 1 at $i = 10$; the plot on the right does not quite.

9.2. **Gromov norm.** Let $G$ be a semigroup acting linearly on a vector space $\mathbf{C}$, and preserving a subspace $\mathbf{B}$. There is a quotient action of $G$ on $\mathbf{H} := \mathbf{C}/\mathbf{B}$. If $\alpha \in \mathbf{H}$ and $G_\alpha$ is the stabilizer of $\alpha$, then there is an affine action of $G_\alpha$ on $\alpha + \mathbf{B}$. This action has a global fixed point only if there is some $a \in \mathbf{C}$ in the class of $\alpha$ fixed by $G_\alpha$.

The sequence $\mathbf{B} \to \mathbf{C} \to \mathbf{H}$ of $G_\alpha$-modules gives rise to a long exact sequence in cohomology:

$$H^0(G_\alpha;\mathbf{B}) \to H^0(G_\alpha;\mathbf{C}) \to H^0(G_\alpha;\mathbf{H}) \to H^1(G_\alpha;\mathbf{B}) \to H^1(G_\alpha;\mathbf{C}) \to \cdots$$

For any $G_\alpha$-module $V$, the group $H^0(G_\alpha;V)$ is equal to $V^{G_\alpha}$, the subspace of $V$ fixed pointwise by $G_\alpha$. There is a tautological class $\alpha \in H^0(G_\alpha;\mathbf{H})$, whose image in $H^1(G_\alpha;\mathbf{B})$ is nontrivial unless it comes from some invariant $a \in H^0(G_\alpha;\mathbf{C})$. Cocycles $c_a$ representing the class in $H^1(G_\alpha;\mathbf{B})$ are in bijection with elements $a \in \mathbf{C}$ representing $\alpha$, by the formula $c_a(g) = ga - a$. Note that this class becomes trivial in $H^1(G_\alpha;\mathbf{C})$.

Now let's specialize to the case that $G = \mathrm{End}(F)$ for $F$ a free group. We can take $\mathbf{C}, \mathbf{B}, \mathbf{H}$ to be $C/\epsilon, B/\epsilon, H_1$ as in § 2. If we choose some nontrivial $\alpha \in H_1(F)$ and let $\mathrm{End}_\alpha$ denote the subsemigroup fixing the class $\alpha$, then for each $\phi \in \mathrm{End}_\alpha$ there is a natural isomorphism between $H_2(F*_\phi)$ and $H_1(F)^\phi$, the subspace of $H_1(F)$ fixed by $\phi$.

*Remark* 9.1. Since $\phi_*$ is represented (with respect to some basis of $H_1(F;\mathbb{Z})$) by an integer matrix, the subspace of $H_1(F;\mathbb{R})$ fixed by $\phi_*$ is defined over $\mathbb{Q}$.

If there is a (pseudo)-norm $\|\cdot\|$ on $\mathbf{B}$ invariant under $G$, then the action of $G_\alpha$ on $\alpha + \mathbf{B}$ is an affine isometric action. Now, the action of $\mathrm{End}_\alpha(F)$ on $B/\epsilon$ with the scl norm is 1-Lipschitz in general, but in fact for *random* endomorphisms $\phi$ the action is actually *isometric*; see [13], Thm. 3.16.

*Remark* 9.2. The Random Isometry Theorem 3.16 from [13] actually proves that the probability that a random endomorphism of length $n$ is an isometry goes to 1 like $1 - C^{-n}$. For a given integral homology class $\alpha \in H_1(F)$ the probability that $\phi \in \mathrm{End}(F)$ preserves $\alpha$ is of order $n^{-k/2}$ where $k$ is the rank of $F$. Thus, even conditioned on $\phi$ lying in $\mathrm{End}_\alpha(F)$, the probability that $\phi$ is isometric goes to 1 exponentially fast in $n$.

The *translation length* $\tau(\phi)$ of $\phi \in \mathrm{End}_\alpha(F)$ on $B/\epsilon$ is equal to the infimum of $\mathrm{scl}(c - \phi(c))$ over all representatives $c$ of the class $\alpha$. By properties of translation length, we may also express this as a limit

$$\tau(\phi) = \lim_{n\to\infty} \frac{1}{n} \mathrm{scl}(c - \phi^n(c))$$

for any $c \in B/\epsilon$ representing $\alpha$.

We now express this translation length in the language of so-called *Gromov norms.*

**Definition 9.3.** Let $X$ be a topological space, and let $A \in H_*(X;\mathbb{R})$. The *Gromov norm* of $A$, denoted $\|A\|$, is the infimum

$$\|A\| := \inf_{[\sum t_i \sigma_i]=A} \sum_i |t_i|$$

where the infimum is taken over all (singular) cycles $\sum t_i\sigma_i$ representing the homology class $A$.

Now, $H_*(X;\mathbb{R}) = H_*(X;\mathbb{Z}) \otimes \mathbb{R}$. In dimension 2, $H_2(X;\mathbb{Z})$ is generated by maps from closed, oriented surfaces $S$ into $X$, and $H_2(\pi_1(X);\mathbb{Z})$ is obtained by quotienting out the image of $\pi_2(X)$ under the Hurewicz map.

**Lemma 9.4** (See e.g. [8], p. 11)**.** *For $A \in H_2(\pi_1(X);\mathbb{Z})$ there is an equality*

$$\|A\|_1 = \inf \frac{-2\chi(S)}{n}$$

*where the infimum is taken over all positive integers $n$, all closed oriented surfaces $S$ without sphere components, and all maps $f : S \to X$ representing $nA$ in $H_2(\pi_1(X);\mathbb{Z})$.*

With this definition we have the following lemma:

**Lemma 9.5** (Gromov norm is translation length)**.** *Let $A \in H_2(F*_\phi)$ be the class corresponding to $\alpha \in H_1(F)^\phi$. There is an equality $\|A\| = 4\tau(\phi)$ where $\|\cdot\|$ denotes the Gromov norm.*

*Proof.* Let $f : S \to K(F*_\phi, 1)$ represent $nA$ for some $n$. We think of $K(F, 1)$ as a horizontal slice of $K(F*_\phi, 1)$ as in § 8.1, and put $f(S) \cap K(F, 1)$ in general position. Then $f(S) \cap K(F, 1)$ is a 1-cycle $c$ representing $\alpha$ and we obtain an inequality in one direction. The inequality in the other direction is obtained by finding a surface $f : S \to K(F, 1)$ with $\partial S$ representing some multiple of $c - \phi(c)$, and gluing this up in the suspension to a closed surface representing a multiple of $A$. □

A surface $f : S \to X$ representing $nA$ is *extremal* if $-2\chi(S)/n = \|A\|_1$. Extremal surfaces do not necessarily exist, but if they do they are injective:

**Lemma 9.6.** *Extremal surfaces are $\pi_1$-injective.*

*Proof.* If $f : S \to X$ is not injective there is a finite index cover $S'$ of $S$ with a simple loop in the kernel (this existence of such a cover follows from LERF for surface groups; see [34]). Then $f' : S' \to X$ can be compressed along this simple loop, producing $f'' : S'' \to X$ with strictly lower norm. Contrapositively, if $S$ is extremal, no such virtual compression is possible. □

9.3. **Extremal surfaces and counting quasimorphisms.** Now let $T_\ell$ denote the verbal traintrack of length $\ell$. Recall from Definition 3.3 the linear operators $\partial : \mathbb{R}[E] \to \mathbb{R}[V]$ and $H : \mathbb{R}[E] \to H_1$, and the subspace $W_\ell = \ker \partial \oplus H$. Define $\widehat{W_\ell} = \ker \partial$ and observe that there is a short exact sequence

$$W_\ell \to \widehat{W_\ell} \to H_1$$

By Lemma 7.5 a random endomorphism of length $n$ has a well-defined action on $W_\ell$, with probability $1 - O(C^{-n})$. The same argument in fact gives an action on $\widehat{W_\ell}$. Again, conditioning on fixing some vector $\alpha \in H_1$, a random endomorphism in $\mathrm{End}_\alpha(F)$ has a well-defined linear action on $\widehat{W_\ell}$ and we obtain therefore a well-defined *affine* linear action on $W_\ell$. Since $B/\epsilon \to W_\ell$ is 1-Lipschitz for scl, it follows that the translation length $\tau_\ell(\phi)$ of $\phi$ on $W_\ell$ (for the affine action) is a lower bound on $\tau(\phi)$.

Recall Theorem 7.6, which says that for and $\ell$ there is a fixed $m$ so that for $\phi \in \mathrm{End}(F)$ a random endomorphism of length $n$, with probability $1 - O(C^{-n})$ the map $\phi : \phi^m(W_\ell) \to \phi^m(W_\ell)$ is an isometry. In fact, the subspace on which it acts isometrically is dual to the family of tripod hyperplanes permuted by $\phi$. Now, conditioning on $\phi$ fixing $\alpha$, the conclusion of the theorem still holds with probability $1 - O(C^{-n})$ (at the cost of adjusting the constant $C$). Because the unit ball in the norm on the fixed subspace is polyhedral, there is an $m$ depending only on $\ell$ so that for random $\phi \in \mathrm{End}(F)$, the map $\phi^m : W_\ell \to W_\ell$ is *projection* to one of finitely many possible fixed subspaces, and therefore random elements of $\mathrm{End}_\alpha(F)$ can be partitioned (nonuniquely) into finitely many subsemigroups $\mathrm{End}_{\alpha,i}$, each of which maps to an affine group of the form $\mathbb{R}^N \rtimes G$ where $G$ is *finite*. There is thus a subsemigroup of finite index in $\mathrm{End}_{\alpha,i}(F)$ whose image in $G$ is finite, and which maps homomorphically to the translation group $\mathbb{R}^N$. As we explain in the next subsection, this homomorphism generalizes one of the summands of the *Johnson homomorphism*.

In any case, for each random $\phi \in \mathrm{End}_\alpha(F)$ the power $\phi^m$ acts affinely on $W_\ell$ by the combination of projection to some subspace, followed by translation of that subspace, realizing the translation length on some further subspace. Dual to this subspace is a fixed subspace of $Q_\ell/H^1$, and the suspension of any class in this subspace defines an element of bounded cohomology $H^2_b(F*_\phi)$ positive on $A$.

9.4. **Euler class and rotation quasimorphism.** Throughout this section we fix a free group $F$, and an element $\alpha \in H_1(F)$. Let $F \to \pi_1(\Sigma)$ be a realization of $F$ as the fundamental group of a compact, oriented surface $\Sigma$, which we suppose to have a hyperbolic structure with geodesic boundary.

Let $\mathrm{End}_{\alpha,\Sigma}(F)$ denote the semigroup of outer endomorphisms of $F$ fixing $\alpha$, and geometric with respect to the realization $\Sigma$. Thus elements of $\mathrm{End}_{\alpha,\Sigma}(F)$ are represented by immersions with geodesic boundary of $\Sigma$ into itself, fixing the homology class $\alpha$.

**Proposition 9.7.** *Let* $\alpha \in H_1(F;\mathbb{Z})$ *be primitive. There is a homomorphism* $\rho_{\alpha,\Sigma} : \mathrm{End}_{\alpha,\Sigma}(F) \to \mathbb{Z}$ *defined as follows. If* $c$ *is any cycle representing* $\alpha$*, and* $\mathrm{rot}_\Sigma$ *is the rotation quasimorphism associated to the realization of* $F$ *as* $\pi_1(\Sigma)$*, then*

$$\rho_{\alpha,\Sigma}(\phi) = \mathrm{rot}_\Sigma(c - \phi(c))$$

*Proof.* Since elements of $\pi_1(\Sigma)$ fix points in $S^1_\infty$, it follows that $\mathrm{rot}_\Sigma$ takes values in $\mathbb{Z}$ (on integral chains).

By the linearity of $\text{rot}_\Sigma$, to show that $\rho$ is a homomorphism it suffices to show that it is well-defined; i.e. that $\text{rot}_\Sigma(c - \phi(c))$ does not depend on the choice of a representative $c$ for the homology class $\alpha$. If we choose a different representative $c'$ then we need to show $\text{rot}_\Sigma(c - c') = \text{rot}_\Sigma(\phi(c - c'))$. But $\text{rot}_\Sigma \circ \phi = \phi^* \text{rot}_\Sigma = \text{rot}_\Sigma$, since $\text{rot}_\Sigma$ depends only on the cyclic order on the ends of $F$ induced by the realization, and any geodesic immersion preserves the cyclic order. $\square$

Note that if we fix $\phi$ and vary $\alpha$, the function $\rho_{\cdot,\Sigma}(\phi)$ defines a homomorphism from $H_1(\Sigma)^\phi$ to $\mathbb{Z}$. Identifying $H_1(\Sigma)^\phi$ with $H_2(F*_\phi)$, we obtain in this way an element of $H^2(F*_\phi)$.

**Definition 9.8.** The cohomology class in $H^2(F*_\phi)$ whose value on $A \in H_2(F*_\phi)$ associated to $\alpha \in H_1(F)^\phi$ is $\rho_{\alpha,\Sigma}(\phi)$ is called the *Euler class* associated to the invariant realization $\Sigma$.

An automorphism of $F$ which is geometric with respect to the realization $\Sigma$ is nothing but an element of the mapping class group of $\Sigma$. If we denote by $\text{Mod}_\alpha(\Sigma)$ the subgroup of the mapping class group fixing $\alpha$, we thereby obtain a homomorphism $\rho_\alpha$ to $\mathbb{Z}$. The *Torelli subgroup* $\mathcal{I}(\Sigma)$ is defined to be the subgroup of $\text{Mod}(\Sigma)$ fixing all of $H_1(\Sigma)$. The homomorphisms $\rho_\alpha$ over all $\alpha$ can be put together into a single homomorphism $\rho : \mathcal{I}(\Sigma) \to H^1(\Sigma)$, which is one summand of the well-known Johnson homomorphism. In fact, the relationship of this summand to rotation quasimorphisms (in the language of Chillingworth [15] as discussed in Remark 5.8) can already by found in Johnson's original paper [26].

If $\partial\Sigma$ is empty or has one component, $\rho : \mathcal{I}(\Sigma) \to H^1(\Sigma)$ is surjective. In general, the image is equal to the image of $H_1(\Sigma)$ under contraction (i.e. cap product) with the intersection form $\omega \in \Lambda^2 H^1(\Sigma)$. In geometric terms, $\rho_\alpha$ is nontrivial on $\text{Mod}_\alpha(\Sigma)$ if and only if there is $\beta$ with $\omega(\alpha, \beta) \neq 0$. However, $\rho_\alpha$ is always nontrivial on $\text{End}_{\alpha,\Sigma}(F)$.

*Example* 9.9. Suppose there is a simple, separating geodesic $\gamma$ representing $\alpha$ (for example, $\gamma$ could be a boundary component of $S$ if $\partial S$ has at least 2 components). Let $\gamma'$ be an immersed geodesic homologous to $\gamma$ with $\text{rot}_\Sigma(\gamma - \gamma') = 1$. To see that there is such a $\gamma'$, first take *any* curve homologous to $\gamma$, and insert a suitable power of $[a, b]$ into a cyclic word representing the conjugacy class, where $a, b$ are any elements of $\pi_1(\Sigma)$ generating a subgroup equal to the fundamental group of some immersed once-punctured torus.

Now, there is a cover $\Sigma'$ of $\Sigma$ containing an embedded lift of $\gamma'$, and since $\gamma'$ is immersed, it separates $\Sigma'$ into two nontrivial pieces. There is a further finite cover $\Sigma''$ of $\Sigma'$ to which $\gamma'$ still lifts, where the pieces on either side have arbitrarily large genus. Then the components of $\Sigma - \gamma$ embed into the components of $\Sigma'' - \gamma'$, and composition with the covering projection $\Sigma'' \to \Sigma$ determines an immersion of $\Sigma$ to itself taking $\gamma$ to $\gamma'$.

In a similar way one can show that $\rho_{\alpha,\Sigma}$ always surjects onto $\mathbb{Z}$.

### 9.5. **Flat surfaces.**

**Definition 9.10.** Let $\phi \in \text{End}_{\alpha,\Sigma}(F)$. A *flat surface* is an immersion $S \to \Sigma$ with geodesic boundary, so that $\partial S = \gamma - \phi(\gamma)$ for some (immersed) geodesic 1-manifold $\gamma$ representing $\alpha$.

A flat surface glues up in the suspension to produce a closed surface mapping to $K(F*_\phi, 1)$. By abuse of notation, we also refer to this closed surface as a flat surface.

**Lemma 9.11.** *A flat surface is extremal, and therefore* $\pi_1$*-injective.*

*Proof.* To see that a flat surface is extremal, it suffices (by Lemma 9.5) to see that $\mathrm{scl}(\gamma - \phi(\gamma))$ is the infimum of $\mathrm{scl}(c - \phi(c))$ over all $c$ representing $\alpha$. But this is certified by Bavard duality and the existence of the extremal dual quasimorphism $\mathrm{rot}_\Sigma$. $\square$

The next lemma shows how to find flat surfaces in practice. The point of this lemma is that both sides of the formula can be computed efficiently in practice.

**Lemma 9.12.** *There is a flat surface* $S$ *with* $\partial S = \gamma - \phi(\gamma)$ *if and only if* $\mathrm{scl}(\gamma - \phi(\gamma)) = \frac{1}{2}\mathrm{rot}_\Sigma(\gamma - \phi(\gamma))$.

*Proof.* This is immediate from the definitions. $\square$

A flat surface has Gromov norm zero if and only if it is a torus; this corresponds precisely to the case that $\phi$ preserves a (homologically essential) conjugacy class in $F$. Otherwise the Gromov norm of such a surface will be positive, and therefore $\rho_{\alpha,\Sigma}(\phi)$ must be nonzero. Conversely, providing $\phi$ does not preserve a nontrivial conjugacy class (for example, if $\phi$ is irreducible), the vanishing of $\rho_{\alpha,\Sigma}(\phi)$ is an obstruction to the existence of a flat surface representing the given homology class.

The definition of flat surfaces makes sense even when $\phi$ is an automorphism of $\Sigma$. In this case the suspension $M_\phi := \Sigma \times [0,1]/(x,0) \sim (\phi(x),1)$ is a 3-manifold fibering over the circle. The Euler class $e_\phi$ of the fibration (i.e. the Euler class of the plane field of tangent planes to the surface fibers) agrees with the Euler class as given in Definition 9.8, explaining the name.

**Lemma 9.13.** *Let* $\phi \in \mathrm{Mod}(\Sigma)$*, and let* $M_\phi$ *be the 3-manifold obtained by suspension. A homology class* $A \in H_2(M_\phi)$ *is represented by a flat surface if and only if the homology class* $A$ *projectively intersects the boundary of the fibered face of the Thurston norm ball in* $H_2(M_\phi, \partial M_\phi)$ *associated to* $e_\phi$*.*

*Proof.* Integral classes in the cone on the fibered face correspond precisely to Thurston norm-minimizing surfaces $S$ for which $e_\phi$ is extremal. These are exactly the surfaces which can be isotoped to be transverse to the suspension flow. Such a transverse surface is flat. Conversely, flat surfaces are norm-minimizing, and certified by $e_\phi$. $\square$

Agol [1] showed that for any 3-manifold $M$ whose fundamental group satisfies a technical condition called *residually finite rational soluble* (usually abbreviated to RFRS), for any nontrivial $A \in H_2(M, \partial M)$ there is a finite cover of $M$ for which the transfer of $A$ is projectively in the closure of a fibered face. The RFRS condition is now known (by work of Wise [42]) to hold for all 3-manifold groups with $H_2$ positive; it follows that every norm-minimizing surface is virtually flat.

*Example* 9.14. Let $S$ be a genus 2 surface with one boundary component. Let $T$ be an embedded once-punctured torus, and let $\gamma$ be a nonseparating essential simple closed curve in $S - T$. Form $\gamma'$ by connect summing $\gamma$ to $\partial T$ along an embedded arc disjoint from either. Then $\gamma - \gamma'$ bounds an embedded twice-punctured torus

$T'$. Since $\gamma$ and $\gamma'$ are both nonseparating, there are many $\phi \in \mathrm{Mod}(S)$ taking $\gamma$ to $\gamma'$. For any such $\phi$, the surface $T'$ glues up to a flat surface in the suspension.

*Example* 9.15. Let $F = \langle x, y\rangle$ and $\phi(x) = xYxY$, $\phi(y) = yyXYx$. Then $\phi([y]) = [y]$. Let $\Sigma$ be the basic surface realization given by the cyclic order $xyXY$. One can check that $\phi$ is geometric with respect to $\Sigma$. Another way to show this is to simply check that $\mathrm{rot}_\Sigma(\phi([y, x])) = 1$, since $[y, x]$ is the boundary of this realization, and this shows the image surface is immersed. Finally, we check that $\mathrm{rot}_\Sigma(y - yyXYx) = 1$ and $\mathrm{scl}(y - yyXYx) = 1/2$. By Lemma 9.12, any extremal surface for $y - yyXYx$ is flat. Using `scallop` [14] (made easier by `wallop` [40]), we can find a particular example: a genus 1 surface with boundary $yy + XyxYYXyxYY$. A set of generators for this surface group is $\{yy, Xyyx, XXyx, t\}$, where recall $F*_\phi = \langle x, y, t \,|\, txt^{-1} = \phi(x), tyt^{-1} = \phi(y)\rangle$

*Example* 9.16. Let $F = \langle x, y, z\rangle$, and let $\phi : F \to F$ be defined

$$x \mapsto YZXYz, \quad y \mapsto ZZXzx, \quad z \mapsto zYXzX.$$

Then $\phi$ is geometric with respect to the realization $\Sigma$ given by the cyclic order $acCbAB$, as described above, and $\phi([x] + [ZZ]) = [x] + [ZZ]$. We also have $\mathrm{scl}(x + ZZ + Zyxzy + xZxyZxyX) = 3/2$, and $\mathrm{rot}_\Sigma(x + ZZ + Zyxzy + xZxyZxyX) = 3$. Therefore, any extremal surface for this chain gives a norm-realization surface. For example, there is a genus zero surface (punctured sphere) with 8 boundaries which bounds twice the chain. Gluing this surface produces a closed genus 4 norm-realizing surface in $F*_\phi$.

*Example* 9.17 (Flat branched surface). Suppose $\phi : F \to F$ is geometric with respect to some realization, so that it is realized as an immersion $f : S \to S$. If we fix a hyperbolic structure on $S$, we can insist that $f$ takes $\partial S$ to a collection of geodesics in $S$. The immersed surface $f(S)$ lifts to an embedding in a finite cover $\widehat{S} \to S$, and the difference $S' := \widehat{S} - f(S)$ is an essential surface, immersing in $S$, and bounding $n\partial S - f(\partial S)$ for some positive integer $n$. Some finite cover of the surface $S'$ glues up in the HNN extension to produce an oriented *branched surface* $S''$ with the property that every oriented surface carried by $S''$ is flat (note that such oriented surfaces necessarily have boundary). Now, $\pi_1(S'')$ contains closed surface subgroups, but these surfaces are not oriented compatibly with $S''$. Is it possible that some of these surface subgroups map injectively to $F*_\phi$?

We end our discussion of extremal and flat surfaces with two natural questions.

**Question 9.18.** *Let $F*_\phi$ be an HNN extension associated to an endomorphism $\phi$. Is the unit ball in the Gromov norm on $H_2(F*_\phi)$ a finite-sided rational polyhedron? Is every rational class projectively represented by an extremal surface? Is every extremal surface dual to an extremal quasimorphism obtained from an invariant class in some $Q_\ell$?*

**Question 9.19.** *Let $F*_\phi$ be an HNN extension associated to an endomorphism $\phi$. Is every homology class in $H_2(F*_\phi)$ virtually dual to the Euler class of some virtual invariant realization?*

9.6. **Experimental results.** For concreteness, consider endomorphisms of length at most 5 in a rank 2 free group. There are 25,544 endomorphisms with non-elementary image, satisfying $|\phi_c(x)| \geq 1$ for each generator $x$, and fixing some primitive vector in homology. By Theorem 7.24 these are all virtually geometric. For computational reasons, we focus on the $6,136$ for which $\phi^2$ is geometric. Of these, we find 940 which contain certified flat surface subgroups.

## 10. Acknowledgments

We would like to thank Sang-Hyun Kim and Henry Wilton for helpful conversations about the material in this paper. Danny Calegari was supported by NSF grant DMS 1005246, and Alden Walker was supported by NSF grant DMS 1203888.

## References

[1] I. Agol, *Criteria for virtual fibering*, J. Topol. **1** (2008), no. 2, 269–284

[2] I. Agol, *The virtual Haken conjecture*, preprint arXiv:1204.2810

[3] C. Bavard, *Longueur stable des commutateurs*, Enseign. Math. (2), **37**, 1-2, (1991), 109–150

[4] M. Bestvina and M. Handel, *Train tracks and automorphisms of free groups*, Ann. Math. **135** (no. 1), (1992), 1–51

[5] R. Brooks, *Some remarks on bounded cohomology*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978), pp. 53–63, Ann. of Math. Stud., **97**, Princeton Univ. Press, Princeton, N.J., 1981

[6] D. Calegari, *Foliations and the geometry of* 3*-manifolds*, Oxford Mathematical Monographs. Oxford University Press, Oxford, 2007

[7] D. Calegari, *Surface subgroups from homology*, Geom. Topol. **12** (2008), no. 4, 1995–2007

[8] D. Calegari, *scl*, MSJ Memoirs, **20**. Mathematical Society of Japan, Tokyo, 2009.

[9] D. Calegari, *Stable commutator length is rational in free groups*, Jour. Amer. Math. Soc. **22** (2009), no. 4, 941–961

[10] D. Calegari, *Faces of the scl norm ball* Geom. Topol. **13** (2009), 1313–1336

[11] D. Calegari, *Scl, sails and surgery*, J. Topol. **4** (2011), no. 2, 305–326

[12] D. Calegari and A. Walker *Random rigidity in the free group*, preprint; arXiv:1104.1768.

[13] D. Calegari and A. Walker *Isometric endomorphisms of free groups*, New York Journal of Math, **17** (2011) 713–743

[14] D. Calegari and A. Walker, `scallop`, computer program available from the authors' webpages, and from `computop.org`

[15] D. Chillingworth, *Winding numbers on surfaces. I.*, Math. Ann. **196** (1972), 218–249

[16] J. Crisp, M. Sageev and M. Sapir, *Surface subgroups of right-angled Artin groups*, Internat. J. Algebra Comput. **18** (2008), no. 3, 443–491

[17] M. Culler, *Using surfaces to solve equations in free groups*, Topology **20** (1981), no. 2, 133–145

[18] N. Dunfield and W. Thurston, *The virtual Haken conjecture: experiments and examples*, Geom. Topol. **7** (2003), 399–441

[19] D. Epstein and K. Fujiwara, *The second bounded cohomology of word-hyperbolic groups*, Topology **36** (1997), no. 6, 1275–1289

[20] GNU Linear Programming Kit, Version 4.45,
`http://www.gnu.org/software/glpk/glpk.html`

[21] C. Gordon, D. Long and A. Reid, *Surface subgroups of Coxeter and Artin groups*, J. Pure Appl. Algebra **189** (2004), no. 1–3, 135–148

[22] C. Gordon and H. Wilton, *On surface subgroups of doubles of free groups*, J. Lond. Math. Soc. (2) **82** (2010), no. 1, 17–31

[23] R. Grigorchuk, *Some results on bounded cohomology*, Combinatorial and geometric group theory (Edinburgh, 1993), 111–163 LMS Lecture Note Ser. **204**, Cambridge Univ. Press, Cambridge, 1995

[24] M. Gromov, *Volume and bounded cohomology*, Inst. Hautes Études Sci. Publ. Math. (1982), no. 56, 5–99

[25] Gurobi Optimization, Inc., *Gurobi Optimizer Reference Manual* (2012),
`http://www.gurobi.com`

[26] D. Johnson, *An abelian quotient of the mapping class group* $\mathcal{I}_g$, Math. Ann. **249** (1980), no. 3, 225–242

[27] J. Kahn and V. Markovic, *Immersing almost geodesic surfaces in a closed hyperbolic three manifold*, Ann. of Math. (2) **175** (2012), no. 3, 1127–1190

[28] S.-H. Kim and S.-I. Oum, *Hyperbolic surface subgroups of one-ended doubles of free groups*, preprint; arXiv:1009.3820

[29] S.-H. Kim and H. Wilton, *Polygonal words in free groups*, Q. J. Math. **63** (2012), no. 2, 399–421

[30] R. Penner, *Perturbative series and the moduli space of Riemann surfaces*, J. Diff. Geom. **27** (1988), 35–53

[31] P. Reynolds, *Dynamics of Irreducible Endomorphisms of* $F_n$, preprint; arXiv:1008.3659

[32] A. Rhemtulla, *A problem of bounded expressibility in free products*, Proc. Cambridge Philos. Soc. **64** (1968), 573–584

[33] M. Sapir, *Some group theory problems*, Internat. J. Algebra Comput. **17** (2007), no. 5–6, 1189–1214

[34] P. Scott, *Subgroups of surface groups are almost geometric*, J. Lond. Math. Soc. **17** (1978), no. 3, 555–565

[35] R. Sharp, *Local limit theorems for free groups*, Math. Ann. **321** (2001), no. 4, 889–904

[36] J. Stallings, *Topology of finite graphs*, Invent. Math. **71** (1983), no. 3, 551–565

[37] W. A. Stein et al., Sage Mathematics Software (Version 5.3), The Sage Development Team (2012) `http://www.sagemath.org`.

[38] W. Thurston, *Geometry and Topology of 3-manifolds*, Notes from lecture at Princeton University, available from the MSRI (a.k.a. "Thurston's Notes")

[39] A. Walker, `gallop`, computer program available from the author's webpage

[40] A. Walker, `wallop`, computer program available from the author's webpage

[41] H. Wilton, *One-ended subgroups of graphs of free groups with cyclic edge groups* Geom. Topol. **16** (2012), 665–683

[42] D. Wise, *The structure of groups with a quasiconvex hierarchy*, preprint; available from the author's webpage

University of Chicago, Chicago, Ill 60637 USA
*E-mail address*: `dannyc@math.uchicago.edu`

University of Chicago, Chicago, Ill 60637 USA
*E-mail address*: `akwalker@math.uchicago.edu`